

Perfect kernel and dynamics: from Bass-Serre theory to hyperbolic groups

P en lope Azuelos and Damien Gaboriau

October 3, 2023

Abstract

We introduce several approaches to studying the Cantor-Bendixson decomposition of and the dynamics on the (topological) space of subgroups for various families of countable groups. In particular, we uncover the perfect kernel and the Cantor-Bendixson rank of the space of subgroups of many new groups, including for instance infinitely ended groups, limit groups, hyperbolic 3-manifold groups and many graphs of groups. We also study the topological dynamics of the conjugation action on the perfect kernel, establishing the conditions for topological transitivity and higher topological transitivity. As an application, we obtain many new examples of groups in the class \mathcal{A} of Glasner and Monod, i.e. admitting faithful transitive amenable actions. This includes for example right-angled Artin groups, limit groups, $C'(1/6)$ small cancellation groups, random groups at density $d < 1/6$, and more generally all virtually compact special groups.

Keywords: Space of subgroups; perfect kernel; topologically transitive actions; hyperbolic groups, subgroup separable groups, Bass-Serre theory.

MSC-classification: 37B; 20E; 20F67; 20F65; 37B20

Contents

1	Introduction	2
2	Notation and preliminaries	8
3	Generalities on the topological structure of $\text{Sub}(G)$	9
3.1	Normal subgroups and invariant subsets	9
3.2	Examples	12
3.3	Subgroups with Schreier graph quasi-isometric to \mathbb{Z}	13
4	Subgroup separable groups and the first ℓ^2-Betti number	16
5	Actions on trees	19
5.1	Main technical dendrological lemma	21
5.2	Main dendrological theorems	24
5.3	Applications	30
5.4	A rich class of groups with maximal perfect kernel	32

6	Hyperbolic groups	35
6.1	Generalities on the space of subgroups of hyperbolic groups	35
6.2	Proof of Theorems 6.1 and 6.2	36
6.3	Hyperbolic 3-manifold groups	41
7	Some facts and more questions	43
8	An application: amenable faithful transitive actions	44
A	Appendix: A list of examples	49

1 Introduction

Let G be a countable group and $\text{Sub}(G)$ be the space of all subgroups of G . There has been a recent increase in studies of the action by conjugation of G on this space. It is the crucible where certain properties of the non-free actions of G boil down, whether they are of a topological or measured nature.

When endowed with the Chabauty topology, $\text{Sub}(G)$ is a compact metrizable totally disconnected space, thus satisfying the Baire property. The Cantor–Bendixson structure theory of compact spaces leads us to consider the unique decomposition $\text{Sub}(G) = \mathcal{K}(G) \sqcup C$ where $\mathcal{K}(G)$, the **perfect kernel** of G , is the maximal perfect subspace of $\text{Sub}(G)$ and C is a countable set. Associated to this decomposition is an ordinal, called the **Cantor-Bendixson rank** of G and denoted $\text{rk}_{\text{CB}}(G)$. It captures after how many steps the iterative process of removing isolated points from $\text{Sub}(G)$ results in the stable subspace $\mathcal{K}(G)$. Precise definitions are given in Section 2.

The action of G on its space of subgroups by conjugation is continuous and the perfect kernel is invariant under this action. The interesting uniformly recurrent subgroups (URS) and invariant random subgroups (IRS) are supported on the perfect kernel.

It is an instructive exercise to check that $\mathcal{K}(\mathbb{Z}^d) = \emptyset$ and that $\text{rk}_{\text{CB}}(\mathbb{Z}^d) = d + 1$. In fact, since a non-empty perfect Polish space contains a Cantor set, $\mathcal{K}(G) = \emptyset$ if and only if G admits only countably many subgroups. This is the case for instance for all finitely generated nilpotent groups. More generally, this happens for all groups whose subgroups are all finitely generated (the so-called Noetherian groups), for instance for the Tarski Monsters: those non-abelian groups whose proper subgroups are all cyclic.

If G is the \mathfrak{R} -free group of countably infinite rank of a (non-trivial) variety \mathfrak{R} of groups (such as the free group \mathbf{F}_∞ , $\oplus_{\mathbb{N}}\mathbb{Z}$, $\oplus_{\mathbb{N}}\mathbb{Z}/m\mathbb{Z}$, the free Burnside group $B(\mathbb{N}, p) = \langle (s_i)_{i \in \mathbb{N}} \mid g^p = 1, \text{ for all } g \rangle$ for $p \geq 2, \dots$), then $\mathcal{K}(G) = \text{Sub}(G)$ (see Proposition 3.8) and $\text{rk}_{\text{CB}}(G) = 0$.

In general, if G is finitely generated then its finite index subgroups are isolated, so

$$\mathcal{K}(G) \subseteq \text{Sub}_{[\infty]}(G) \text{ where } \text{Sub}_{[\infty]}(G) := \{H \leq G \text{ s.t. } [G : H] = \infty\}.$$

In the case where $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$, note that $\text{rk}_{\text{CB}}(G) = 1$. Observe that $\text{Sub}(G)$ may contain infinite index isolated points. There are even examples of infinite groups for which the trivial subgroup $\{\text{id}\}$ is isolated: consider for instance the Prüfer p -group, or see the more interesting finitely generated Example 3.7.

It is not hard to check that the equality holds for all finitely generated non-cyclic free groups \mathbf{F}_r :

$$\mathcal{K}(\mathbf{F}_r) = \text{Sub}_{[\infty]}(\mathbf{F}_r),$$

see e.g. [CGLM20, Proposition 2.1]. Moreover, the action of \mathbf{F}_r on $\mathcal{K}(\mathbf{F}_r)$ is highly topologically transitive (see Corollary 5.17 or Corollary 6.3). Recall that a continuous action $G \curvearrowright X$ on a compact metrizable space of a countable group G is called **topologically r -transitive** for a positive integer r if the diagonal action $G \curvearrowright X^r$ on the r -fold space is **topologically transitive**; i.e. if for every $2r$ -tuple $(V_1, V_2, \dots, V_{2r})$ of non-empty open sets, there is an element $g \in G$ such that $g \cdot V_i \cap V_{r+i} \neq \emptyset$ for all $i = 1, 2, \dots, r$ (see Section 2). The action is called **highly topologically transitive** if it is topologically r -transitive for all $r \geq 1$.

One purpose of this article is to extend the validity of these statements to several new groups.

In recent years, the space of subgroups of several classes of groups have been studied by various authors. Cornulier, Guyot and Pitsch [CGP10] classified the spaces of subgroups of all countable abelian groups up to homeomorphism and computed their Cantor-Bendixson rank. In [BGK15], Bowen, Grigorchuk and Kravchenko found the perfect kernel and Cantor-Bendixson rank of the lamplighter groups $(\mathbb{Z}/k\mathbb{Z})^n \wr \mathbb{Z} = \bigoplus_{\mathbb{Z}} (\mathbb{Z}/k\mathbb{Z})^n \rtimes \mathbb{Z}$, i.e. $\mathcal{K}((\mathbb{Z}/k\mathbb{Z})^n \wr \mathbb{Z}) = \text{Sub}(\bigoplus_{\mathbb{Z}} (\mathbb{Z}/k\mathbb{Z})^n)$ and $\text{rk}_{\text{CB}}((\mathbb{Z}/k\mathbb{Z})^n \wr \mathbb{Z}) = \omega$, the first infinite ordinal. Skipper and Wesolek described in [SW20] the perfect kernels of the Grigorchuk group, the Gupta-Sidki 3-group and some other families of groups of automorphisms of infinite regular rooted trees; they also established that $\text{rk}_{\text{CB}}(G) = \omega$ for these groups.

Recently Carderi, Le Maître, Stalder and the second author described the perfect kernel and the dynamics for the Baumslag-Solitar groups $\text{BS}(m, n) = \langle b, t | tb^m t^{-1} = b^n \rangle$ depending on the parameters m, n in [CGLMS22]. If $|m|, |n| > 1$, then $\mathcal{K}(\text{BS}(m, n))$ is the space of subgroups whose action on the standard Bass-Serre tree of $\text{BS}(m, n)$ has infinite quotient. Furthermore, the perfect kernel admits a countably infinite partition into $\text{BS}(m, n)$ -invariant subspaces. One of them is closed and not open; all the others are open (and also closed exactly when $|m| = |n|$). Moreover, the $\text{BS}(m, n)$ -action on each piece is topologically transitive.

Intrinsically, this theory is less interested in the algebraic structure of subgroups so much as the “geometry of their embedding” into G . A result which exemplifies this is the following: *If $H \leq G$ are finitely generated groups and the quotient $H \backslash G$ is quasi-isometric to \mathbb{Z} then H vanishes by the third Cantor-Bendixson derivative of $\text{Sub}(G)$* (Theorem 3.10).

Hyperbolic groups An important class of finitely generated groups is that of hyperbolic groups, introduced by Gromov [Gro87]. Describing the space of subgroups of a hyperbolic group is in general a difficult task, largely due to the fact that finitely generated subgroups of hyperbolic groups need not respect the geometry of the group. There are however a range of tools available to study those subgroups that do respect the geometry: namely the quasiconvex subgroups.

Given a hyperbolic group G , we denote by $\text{Sub}_{\text{QC}[\infty]}(G)$ the set of quasiconvex subgroups of G with infinite index.

Theorem A (Theorems 6.1 and 6.2). *Let G be a non-elementary hyperbolic group. Then*

$$\overline{\text{Sub}_{\text{QC}[\infty]}(G)} \subseteq \mathcal{K}(G).$$

If, in addition, the maximal finite normal subgroup of G is trivial then the action of G on $\overline{\text{Sub}_{\text{QC}[\infty]}(G)}$ is highly topologically transitive.

A hyperbolic group is said to be locally quasiconvex if each of its finitely generated subgroups is quasiconvex. There are many examples of hyperbolic groups which satisfy this property, including (but not limited to): hyperbolic surface groups, hyperbolic limit groups, certain one-relator groups with torsion, certain small cancellation groups and graphs of free groups with cyclic edge groups which do not contain any Baumslag-Solitar subgroups (see Examples 6.4, 6.5, 6.6, and 6.7, and the references therein).

Corollary B (Corollary 6.3). *If a group G is non-elementary hyperbolic and locally quasiconvex then*

$$\mathcal{K}(G) = \text{Sub}_{[\infty]}(G) = \overline{\text{Sub}_{\text{QC}[\infty]}(G)}.$$

If in addition the maximal finite normal subgroup of G is trivial then the action of G on $\mathcal{K}(G)$ is highly topologically transitive.

Another class of hyperbolic groups of particular interest is that of hyperbolic 3-manifold groups. While these are not locally quasiconvex, their finitely generated subgroups are well understood. As a result, we are able to describe their perfect kernels in their entirety.

Theorem C (Theorem 6.12). *Let M be a closed hyperbolic 3-manifold and $G = \pi_1(M)$. Then*

$$\mathcal{K}(G) = \overline{\text{Sub}_{\text{QC}[\infty]}(G)}.$$

More precisely, the perfect kernel of G is the set of infinite index subgroups $H \leq G$ such that either H is quasiconvex or H is not finitely generated.

Moreover, the action of G on $\mathcal{K}(G)$ is highly topologically transitive and the Cantor-Bendixson rank of G is 2 or 3.

The above theorem makes use of significant discoveries which have been made about 3-manifolds over the past few decades. In particular, we use a consequence of the Tameness Theorem [Ago04], [CG06] and Canary's Covering Theorem [Can96] as well as the fact that closed hyperbolic 3-manifolds are virtually fibered [Ago13] (by virtue of being virtually special in the sense of [HW08]).

In general, the inclusion $\overline{\text{Sub}_{\text{QC}[\infty]}(G)} \subseteq \mathcal{K}(G)$ is not an equality. Indeed, Rips' construction [Rip82] allows us to find a hyperbolic group G so that we can embed a G -invariant open subset U into the complement $\mathcal{K}(G) \setminus \overline{\text{Sub}_{\text{QC}[\infty]}(G)}$. Moreover, one can ensure that the action of G on U is as pathological as that of any finitely presented group on its perfect kernel. We discuss this in more detail in Section 7.

Subgroup separability A group G is **subgroup separable** (also referred to as LERF or locally extended residually finite) if every finitely generated subgroup of G can be expressed as an intersection of finite index subgroups. The class of groups with this property includes free groups [Hal49] and surface groups [Sco78].

Subgroup separability is equivalent to the density in $\text{Sub}(G)$ of the set of finite index subgroups, as observed in [GKM16] (see Section 4). This does not provide any further information on the perfect kernel of a group, except when we can relate the rank and the index of finite index subgroups. For this purpose, we require that the first ℓ^2 -Betti number be positive, we will see that in many cases this condition is in fact optimal (see Remark 4.2).

Theorem D (Theorem 4.1). *Let G be a countable and subgroup separable group. If G is not finitely generated then $\text{Sub}(G)$ is a perfect space. If G is finitely generated with positive first ℓ^2 -Betti number then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ and $\text{rk}_{\text{CB}}(G) = 1$.*

As mentioned above, free groups and surface groups are subgroup separable. Moreover, hyperbolic surface groups have positive first ℓ^2 -Betti number, as do non-abelian free groups. Thus the above theorem implies that the perfect kernel of a finite rank non-abelian free group or a hyperbolic compact surface group coincides with its set of infinite index subgroups, and its Cantor-Bendixson rank is 1. Note that the remaining (non-hyperbolic) compact surface groups are virtually abelian so their perfect kernel is empty.

More generally, we have the following corollary for Sela's limit groups. Recall that a group G is a **limit group**, or **ω -residually free**, if for any finite set $S \subseteq G \setminus \{\text{id}\}$ there exists a normal subgroup $N \trianglelefteq G$ such that the quotient G/N is free and $S \cap N = \emptyset$. Equivalently, limit groups are the limits of free groups in the space of marked groups [CG05, Theorem 1.1].

Corollary E (Limit groups (Corollaries 4.4 and 5.23)). *If G is a finitely generated non-abelian limit group, then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ and the action $G \curvearrowright \mathcal{K}(G)$ is highly topologically transitive.*

Groups with infinitely many ends and actions on trees A fundamental notion in geometric group theory is the notion of ends of a group. Recall that a finitely generated group has either zero, one, two or infinitely many ends. Stallings' Theorem [Sta68, Sta71] states that a group has infinitely many ends if and only if it acts non-trivially and irreducibly on a tree with finite edge stabilisers. This theorem is invaluable in the study of infinitely ended groups and, in the context of spaces of subgroups, it allows us to show the following.

Theorem F (Infinitely many ends (Corollary 5.17)). *If G is a finitely generated group with infinitely many ends, then the perfect kernel of G consists of its subgroups with infinite index:*

$$\mathcal{K}(G) = \text{Sub}_{[\infty]}(G).$$

Moreover, if G has no non-trivial finite normal subgroups then the action of G on $\mathcal{K}(G)$ is highly topologically transitive.

It is also interesting to consider actions on trees where stabilisers of individual edges are not necessarily finite or trivial but rather where the stabilisers of sufficiently long paths are trivial. These actions are known as **acylindrical**. Examples of groups admitting such actions include amalgamated free products $A *_C B$ where C is malnormal in A or B (See Section 5.3).

Corollary G (Acylindrical actions (see Corollary 5.15)). *Let $G \curvearrowright T$ be a minimal irreducible and acylindrical action of a countable group on a tree. If $H \leq G$ is a subgroup such that the quotient graph $H \backslash T$ is infinite then $H \in \mathcal{K}(G)$.*

An acylindrical action on a tree only provides information on those subgroups whose quotient is infinite. We construct examples of finitely generated groups G which act acylindrically on trees and which contain infinite index subgroups H whose quotients are finite. Sometimes these subgroups H are in the perfect kernel and sometimes they are not (see Remark 5.16). More information about the group G is therefore required to describe the perfect kernel fully.

The conclusion of Corollary [G](#) in fact holds under a weaker condition than acylindricity of the action. Given a group G acting on a tree T , define the set

$$\text{Sub}_{|\bullet \setminus T|_\infty}(G) := \{H \in \text{Sub}(G) : |H \setminus T| = \infty\},$$

of subgroups that act on T with infinite quotient graph.

Theorem H (Theorems [5.9](#) and [5.11](#)). *Assume $G \curvearrowright T$ is a minimal and irreducible action on a tree. Assume that there are two edges f_1, f_2 of T such that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite. Then*

$$\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} \subseteq \mathcal{K}(G).$$

If, in addition, the kernel of the action $G \curvearrowright T$ is trivial then the action of G on $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$ is highly topologically transitive.

This result allows us to establish that $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ for a large family of groups (namely the class \mathcal{Q} , see Definition [5.21](#)), which includes graphs of free groups with cyclic edge groups such that at least one vertex group is non-cyclic. It also allows us to deduce the topological transitivity properties of the action $G \curvearrowright \mathcal{K}(G)$ (see Corollary [5.23](#)).

The reader will probably have noticed that our results above concerning the dynamics of the action $G \curvearrowright \mathcal{K}(G)$ exclude groups G that contain a non-trivial finite normal subgroup N_0 . They may have wondered what happens in this case. In fact, the action of G on $\text{Sub}(N_0)$ by conjugation induces a finite G -invariant clopen partition of $\text{Sub}(G)$ and of $\mathcal{K}(G)$ (see Proposition [3.2](#)). When we allow the existence of such an N_0 , under the conditions of Theorems [A](#) and [H](#), the action on each piece of this clopen partition is topologically transitive (see Theorem [6.2](#) and Theorem [5.11](#)). However, in general the action on the pieces is not topologically 2-transitive, see Remark [5.13](#).

An application to transitive amenable actions As a by-product of our investigations, we obtain information on transitive actions. Following Glasner-Monod [[GM07](#)], let \mathcal{A} be the class of countable infinite groups that admit a *faithful transitive and amenable* action.

This class contains all free groups and most free products (see [[GM07](#)]), surface groups and hyperbolic 3-manifolds groups (see [[Moo10](#)]) and some instances of amalgamated free products and HNN-extension over finite or amenable groups (see [[Moo11a](#), [Moo11b](#), [Fim14](#)]).

Our results allow us to greatly enlarge the collection of examples of groups in the class \mathcal{A} . This is done by singling out two subgroups $H_1, H_2 \in \text{Sub}_{[\infty]}(G)$ such that $G \curvearrowright G/H_1$ is faithful while $G \curvearrowright G/H_2$ is amenable (thus isolating two separate conditions: faithfulness and co-amenability) and by finding (thanks to topological transitivity) subgroups whose orbit in $\text{Sub}(G)$ accumulates on both H_1 and H_2 (see Proposition [8.1](#)).

Our Corollary [5.17](#) (on groups with infinitely many ends) admits as a corollary Theorem 3.4 of [[GM07](#)]. The following generalisation was suggested to us by François Le Maître, following the circulation of a previous version of this paper.

Corollary I (of Proposition [8.1](#)). *Assume that the action $G \curvearrowright \text{Sub}_{[\infty]}(G)$ is topologically transitive. Assume that G admits an amenable action with only infinite orbits. Then $G \in \mathcal{A}$.*

Here is a (non-exhaustive) list of groups which we can prove belong to the class \mathcal{A} .

- Any infinitely ended group without a non-trivial finite normal subgroup and with some amenable action with infinite orbits (Corollary 5.17 and Proposition 8.1). For instance, any HNN-extension G of any infinite group H above a pair of finite subgroups with trivial intersection.
- The fundamental group of any non-tree graph of groups where some edge group is malnormal in an incident vertex group (See Corollary 8.4).
- Virtually compact special groups (see Theorem 8.6). In particular:
 - Right-angled Artin groups;
 - $C'(1/6)$ small cancellation groups;
 - Right-angled and hyperbolic Coxeter groups;
 - One-relator groups with torsion;
 - Limit groups;
 - Random groups at density $d < 1/6$;
 - Hyperbolic-by-cyclic groups which are themselves hyperbolic.
- Groups in the class \mathcal{Q}' (Definition 5.21) that admit an infinite index co-amenable subgroup (for instance if the defining graph of groups is not a tree) (Corollary 5.23 and Proposition 8.1). In particular, finite graphs of free groups with cyclic edge groups such that at least one vertex group is non-cyclic (Corollary 8.2).

Acknowledgments We are extremely grateful to Mark Hagen for his helpful comments, for many enlightening discussions on hyperbolic groups and for asking us about the space of subgroups of hyperbolic 3-manifold groups. The first author is particularly grateful for his ongoing support.

We would like to thank Ian Biringer for pointing out an example of a hyperbolic 3-manifold group with Cantor-Bendixson rank 3 and for a discussion which motivated us to include and improve upon the results of Section 3.3.

We thank Alessandro Carderi, François Le Maître and Yves Stalder for many enlightening discussions and for drawing our attention to the example of the group G in Example 3.7.

We also thank François Dahmani for several discussions on hyperbolic groups.

We are grateful to Eli Glasner for asking us whether our previous results about topologically transitive actions could be upgraded to weakly mixing actions and for giving us a few basics on the subject. This led us to consider highly topologically transitive actions. This question was asked during the wonderful conference “Measured Group Theory, Stochastic Processes on Groups and Borel Combinatorics” which took place at the CIRM in May 2023.

We are grateful to Matthias Uschold for his comments on a previous version of this paper.

P. A. is supported by a University of Bristol PhD scholarship. She was supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR) during the academic year 2021-2022, when we initiated this project.

D. G. is supported by the C.N.R.S.

2 Notation and preliminaries

Let G be a countable group. The space of subgroups $\text{Sub}(G)$ of G is a subspace of the set $\{0, 1\}^G$, where each subset of G is identified with an element of $\{0, 1\}^G$ via the characteristic map. The Chabauty topology on $\text{Sub}(G)$ is the subspace topology induced by the product topology on $\{0, 1\}^G$. It follows that $\text{Sub}(G)$ is compact, metrisable, separable and totally disconnected. In particular, it satisfies the Baire property and it is a Polish space. The collection of subsets

$$\mathcal{V}_G(\mathcal{I}, \mathcal{O}) := \{H \leq G : \mathcal{I} \subseteq H, H \cap \mathcal{O} = \emptyset\} \subseteq \text{Sub}(G)$$

for all finite subsets $\mathcal{I}, \mathcal{O} \subseteq G$ forms a basis of clopen sets for the Chabauty topology on $\text{Sub}(G)$. When there is no risk of confusion we will write $\mathcal{V}(\mathcal{I}, \mathcal{O})$ instead of $\mathcal{V}_G(\mathcal{I}, \mathcal{O})$. The action of G by conjugation ($g \cdot H = gHg^{-1}$) on $\text{Sub}(G)$ is continuous.

The Cantor-Bendixson decomposition Let X be a Polish space. The Cantor-Bendixson derivatives of X are defined by transfinite induction. The derived set of X is the complement $X' := X \setminus \text{Is}(X)$ of the set $\text{Is}(X)$ of isolated points of X . Let $X^{(0)} = X$. The **Cantor-Bendixson derivatives** of X are defined by repeatedly applying the derived set operation using transfinite recursion: Given an ordinal α such that the α -th Cantor-Bendixson derivative $X^{(\alpha)}$ is defined, let $X^{(\alpha+1)} := (X^{(\alpha)})'$. If β is a limit ordinal such that $X^{(\alpha)}$ is defined for each $\alpha < \beta$, then let $X^{(\beta)} := \bigcap_{\alpha < \beta} X^{(\alpha)}$. The Cantor-Bendixson theorem [Can84, Ben00] implies that there is an, at most countable, ordinal α such that $X^{(\alpha)} = X^{(\alpha+1)}$. In this case, $X^{(\alpha)}$ is the (unique) maximal perfect subspace of X , called the **perfect kernel** of X ; we denote it $\mathcal{K}(X) := X^{(\alpha)}$. The minimal ordinal α such that $X^{(\alpha)} = \mathcal{K}(X)$ is called the **Cantor-Bendixson rank** of X and is denoted by $\text{rk}_{\text{CB}}(X)$. A point $x \in X$ is a **condensation point** if every open neighbourhood of x is uncountable. It follows from the proof of the Cantor-Bendixson theorem (see e.g. [Kec95] for an accessible account) that $x \in \mathcal{K}(X)$ if and only if x is a condensation point.

When G is a countable group, we define

$$\begin{aligned} \mathcal{K}(G) &:= \mathcal{K}(\text{Sub}(G)) \text{ the } \mathbf{perfect\ kernel\ of\ } G, \text{ and} \\ \text{rk}_{\text{CB}}(G) &:= \text{rk}_{\text{CB}}(\text{Sub}(G)) \text{ the } \mathbf{Cantor-Bendixson\ rank\ of\ } G. \end{aligned}$$

Be careful not to confuse this with the Cantor-Bendixson rank $\text{cb}(G)$ that some authors define for a group G as the supremum of the ordinals α such that $\{\text{id}\} \in \mathcal{N}(G)^{(\alpha)}$ (where $\mathcal{N}(G)$ denotes the space of normal subgroups of G) when such an α exists and \mathfrak{C} otherwise, i.e. when $\{\text{id}\} \in \mathcal{K}(\mathcal{N}(G))$ (see for instance [Cor11]).

Let $H \leq G$ be a subgroup. The **Cantor-Bendixson erasing rank** $\text{rk}_{\text{CB}}^{\times}(H; G)$ of H in G is, when $H \notin \mathcal{K}(G)$, the infimum of the ordinals α such that the derivative $\text{Sub}(G)^{(\alpha)}$ does not contain H ; and we write $\text{rk}_{\text{CB}}^{\times}(H; G) = \infty$ when $H \in \mathcal{K}(G)$. We make the convention that ∞ is larger than all the ordinals. Observe that $\text{rk}_{\text{CB}}^{\times}(H; G)$ is not a limit ordinal:

$$\text{rk}_{\text{CB}}^{\times}(H; G) = \begin{cases} \infty & \text{if } H \in \mathcal{K}(G) \\ \alpha + 1 & \text{if } H \in \text{Sub}(G)^{(\alpha)} \text{ but } H \notin \text{Sub}(G)^{(\alpha+1)}. \end{cases}$$

Moreover, $\text{rk}_{\text{CB}}(G) = 0$ if and only if $\text{Sub}(G)$ is perfect and $\text{rk}_{\text{CB}}(G) = \sup\{\text{rk}_{\text{CB}}^{\times}(H; G) : H \in \text{Sub}(G) \setminus \mathcal{K}(G)\}$ otherwise.

Example 2.1. For instance, when $G = \mathbb{Z}^d$, then $\text{rk}_{\text{CB}}^\times(H; \mathbb{Z}^d) = d - \text{rk}(H) + 1$ where $\text{rk}(H)$ is the minimal number of generators of the abelian torsion-free group $H \leq \mathbb{Z}^d$. In particular, $\text{rk}_{\text{CB}}^\times(\{\text{id}\}; \mathbb{Z}^d) = d + 1$, $\text{rk}_{\text{CB}}(\mathbb{Z}^d) = d + 1$, and $\mathcal{K}(\mathbb{Z}^d) = \emptyset$.

Topological transitivity A continuous action $G \curvearrowright X$ of a countable discrete group G on a compact metrizable space is called **topologically transitive** if it satisfies one of the following equivalent conditions:

1. For every pair (V_1, V_2) of non-empty open subsets of X , there is an element $g \in G$ such that $g \cdot V_1 \cap V_2 \neq \emptyset$;
2. There is a dense orbit, i.e. a point $x \in X$ whose orbit $G \cdot x$ is dense in X ;
3. There is a G_δ subset of X consisting of dense orbits.

A continuous action $G \curvearrowright X$ on a compact metrizable space of a countable group G is called **topologically r -transitive** for a positive integer r if the diagonal action $G \curvearrowright X^r$ on the r -fold space is topologically transitive; i.e. if for every $2r$ -tuple $(V_1, V_2, \dots, V_{2r})$ of non-empty open sets, there is an element $g \in G$ such that $g \cdot V_i \cap V_{r+i} \neq \emptyset$ for all $i = 1, 2, \dots, r$. A topologically 2-transitive action is also called **topologically weakly mixing**. If an action is topologically r -transitive for all $r \geq 1$ we say that it is **highly topologically transitive**.

Contrarily to the analogous concept in ergodic theory, topological 2-transitivity does not imply high topological transitivity when the group G is not abelian.

3 Generalities on the topological structure of $\text{Sub}(G)$

In this section we highlight the effects of various properties of individual subgroups, and of their quotients, on the Cantor-Bendixson decomposition of the space of subgroups and the dynamics of the action by conjugation.

Remark 3.1. If $H \leq G$ is a subgroup of the countable group G , then the natural continuous embedding $\text{Sub}(H) \subseteq \text{Sub}(G)$ induces a continuous embedding $\mathcal{K}(H) \subseteq \mathcal{K}(G)$. In fact, the topology on $\text{Sub}(H)$ is induced from that on $\text{Sub}(G)$. Condensation points of $\text{Sub}(H)$ are thus condensation points of $\text{Sub}(G)$.

3.1 Normal subgroups and invariant subsets

A normal subgroup $N \trianglelefteq G$ is clearly a fixed point of the G -action on $\text{Sub}(G)$ by conjugation. In the case where N is finite, this property induces an invariant partition of the entire space.

Proposition 3.2 (Finite normal subgroup and its clopen partition). *If $N \trianglelefteq G$ is a finite normal subgroup N . The set $\text{Conj}(N)^G$ of G -conjugacy classes of subgroups of N induces a finite G -invariant clopen partition:*

$$\text{Sub}(G) = \bigsqcup_{C \in \text{Conj}(N)^G} \{H \in \text{Sub}(G) : H \cap N \in C\}.$$

Proof. By normality and finiteness, G acts by conjugation on $\text{Sub}(N)$ with finite orbits. These orbits form the set $\text{Conj}(N)^G = \{\{\gamma M \gamma^{-1} : \gamma \in G\} : M \in \text{Sub}(N)\}$. Since $\gamma H \gamma^{-1} \cap N = \gamma(H \cap N) \gamma^{-1}$, the G -orbit of $H \cap N \in \text{Sub}(N)$ is an invariant of the G -orbit of $H \in \text{Sub}(G)$. In addition, for $C \in \text{Conj}(N)^G$, the condition $H \cap N \in C$ is a clopen condition in $\text{Sub}(G)$: it defines a union of basic clopen sets: $\{H \in \text{Sub}(G) : H \cap N \in C\} = \cup_{M \in C} \mathcal{V}(M, N \setminus M)$. \square

If G' is a subgroup of the countable group G , then the map

$$P : \begin{pmatrix} \text{Sub}(G) & \longrightarrow & \text{Sub}(G') \\ H & \longmapsto & H \cap G' \end{pmatrix} \quad (1)$$

is continuous: if $\mathcal{V}_{G'}(\mathcal{I}', \mathcal{O}')$ is a basic open set in $\text{Sub}(G')$ (with $\mathcal{I}', \mathcal{O}'$ finite subsets of G'), then $P^{-1}(\mathcal{V}_{G'}(\mathcal{I}', \mathcal{O}')) = \mathcal{V}_G(\mathcal{I}', \mathcal{O}')$.

Proposition 3.3. *Let $G' \leq G$ be a finite index subgroup and $P : \text{Sub}(G) \rightarrow \text{Sub}(G')$ be the continuous map from (1). Then each fiber of P is countable. It follows that the natural embedding $\mathcal{K}(G') \subseteq \mathcal{K}(G) \cap \text{Sub}(G')$ is an equality: $\mathcal{K}(G') = \mathcal{K}(G) \cap \text{Sub}(G')$.*

Proof of Proposition 3.3. Fix a subgroup $G'_0 \leq G'$. Let $k := [G : G']$ and let $g_1, \dots, g_k \in G$ be such that $G = \sqcup_{i=1}^k g_i G'$. If $H \in P^{-1}(G'_0)$ let $I \subseteq \{1, \dots, k\}$ be such that $H \cap g_i G' \neq \emptyset$ if and only if $i \in I$. Then, for each $i \in I$, there is $h_i \in g_i G'$ such that $H \cap g_i G' = h_i G'_0$. Thus $H = \sqcup_{i \in I} h_i G'_0$. It follows that

$$P^{-1}(G'_0) \subseteq \bigcup_{I \subseteq \{1, \dots, k\}} \left\{ \bigsqcup_{i \in I} h_i G'_0 : h_i \in g_i G' \right\}.$$

Thus $P^{-1}(G'_0)$ is contained in a finite union of countable sets and is therefore countable.

To see that $\mathcal{K}(G) \cap \text{Sub}(G') \subseteq \mathcal{K}(G')$, let $H' \in \text{Sub}(G') \setminus \mathcal{K}(G')$. Pick a countable neighbourhood \mathcal{V}' of H' in $\text{Sub}(G')$. Its pre-image $P^{-1}(\mathcal{V}')$ is a countable neighbourhood of H' in $\text{Sub}(G)$ so $H' \notin \mathcal{K}(G)$. \square

Proposition 3.4. *Let $N \trianglelefteq G$ be a finite normal subgroup of the countable group G . Assume G contains a torsion-free subgroup H such that $\{\text{id}\} \in \mathcal{K}(H)$. Then $\text{Sub}(N) \subseteq \mathcal{K}(G)$.*

Proof. The subgroup H acting by conjugation on the finite subgroup N admits a finite index subgroup H' that acts trivially; i.e. H' commutes with N , and since it is torsion-free, it intersects N trivially, thus $K = H' \times N$ injects naturally as a subgroup of G . If $M = \{\text{id}\} \times M$ is a subgroup of $N = \{\text{id}\} \times N$ and $\mathcal{V} = \mathcal{V}_K(M, \mathcal{O})$ is any basic neighbourhood of M in $\text{Sub}(K)$, with $\mathcal{O} = \{(x_1, n_1)(x_2, n_2), \dots, (x_j, n_j)\}$ where $x_i \in H'$ and $n_i \in N$, then $\mathcal{V}' := \mathcal{V}_{H'}(\emptyset, \{x_1, x_2, \dots, x_j\})$ is a basic neighbourhood of $\{\text{id}\}$ in $\text{Sub}(H')$. By Proposition 3.3, $\{\text{id}\} \in \mathcal{K}(H')$. Thus \mathcal{V}' is uncountable and the uncountable set $\{H' \times M : H' \in \mathcal{V}'\}$ is contained in \mathcal{V} . It follows that $M \in \mathcal{K}(K)$, and so $M \in \mathcal{K}(G)$ by Remark 3.1. \square

Let $H \leq G$ be a subgroup. We denote by

$$\text{Sub}_{H \leq}(G) := \{L \leq G : H \leq L\}$$

the closed subspace of $\text{Sub}(G)$ consisting of subgroups that contain H . In the case where H is generated by a finite set S , then $\text{Sub}_{H \leq}(G) = \mathcal{V}(S, \emptyset)$ is a clopen neighbourhood of H in $\text{Sub}(G)$. If H is normal, then the action of G on $\text{Sub}(G)$ by conjugation induces natural actions on the spaces of subgroups $G \curvearrowright \text{Sub}_{H \leq}(G)$ and $G \curvearrowright \text{Sub}(G/H)$.

Proposition 3.5. *Let $1 \rightarrow N \rightarrow G \xrightarrow{\phi} Q \rightarrow 1$ be a short exact sequence.*

1. *Then the bijection*

$$q : \begin{pmatrix} \text{Sub}_{N \leq}(G) & \rightarrow & \text{Sub}(Q) \\ H & \mapsto & H/N \end{pmatrix}$$

is a G -equivariant homeomorphism with respect to the subspace topology on $\text{Sub}_{N \leq}(G)$. In particular,

$$q^{-1}(\mathcal{K}(Q)) \subseteq \mathcal{K}(G). \quad (2)$$

2. *Assume that N is finitely generated. Then $\text{Sub}_{N \leq}(G)$ is a clopen neighbourhood of N in $\text{Sub}(G)$. It follows that the restriction*

$$q_{\upharpoonright} : \mathcal{K}(G) \cap \text{Sub}_{N \leq}(G) \longrightarrow \mathcal{K}(Q) \quad (3)$$

is a G -equivariant homeomorphism.

If $H \in \text{Sub}_{N \leq}(G)$, then the Cantor-Bendixson erasing ranks of H in G and of $q(H) = H/N$ in Q are the same:

$$\text{rk}_{\text{CB}}^{\times}(H; G) = \text{rk}_{\text{CB}}^{\times}(H/N; Q). \quad (4)$$

In particular, if $\text{Sub}(Q)$ is countable, and if $H \in \text{Sub}_{N \leq}(G)$, then

$$\text{rk}_{\text{CB}}^{\times}(H; G) \leq \text{rk}_{\text{CB}}(Q). \quad (5)$$

Proof. Let us show that the bijection $q : \text{Sub}_{N \leq}(G) \rightarrow \text{Sub}(Q)$ is continuous. If $\mathcal{V}_Q(\mathcal{I}, \mathcal{O})$ is a basic clopen set in $\text{Sub}(Q)$, let \mathcal{I}' and \mathcal{O}' be any (bijective!) ϕ -sections in G of the finite sets \mathcal{I} and \mathcal{O} . Then $q^{-1}(\mathcal{V}_Q(\mathcal{I}, \mathcal{O})) = \mathcal{V}_G(\mathcal{I}', \mathcal{O}') \cap \text{Sub}_{N \leq}(G)$. To see that (2) holds, recall that if $H \in q^{-1}(\mathcal{K}(Q))$ then every open neighbourhood of $q(H)$ in $\text{Sub}(Q)$ is uncountable. Since q is a homeomorphism, it follows that every open neighbourhood of H in $\text{Sub}_{N \leq}(G)$ is uncountable and thus so is every open neighbourhood of H in $\text{Sub}(G)$. Hence $H \in \mathcal{K}(G)$.

If N is generated by the finite set S , then $\text{Sub}_{N \leq}(G)$ coincides with the basic clopen set $\mathcal{V}_G(S, \emptyset)$.

If Y is an open subset of a Polish space X then a point $y \in Y$ is isolated with respect to the subspace topology on Y if and only if y is isolated in X . It follows by transfinite induction that, for every $y \in Y$ and every ordinal α , we have $y \in Y^{(\alpha)}$ if and only if $y \in X^{(\alpha)}$. Thus $Y^{(\alpha)} = Y \cap X^{(\alpha)}$. Again by transfinite induction, one can show that, if $f : X \rightarrow Y$ is a homeomorphism between Polish spaces X, Y , then $f(X^{(\alpha)}) = Y^{(\alpha)}$ for all α . In particular, $f(\mathcal{K}(X)) = \mathcal{K}(Y)$ and the Cantor-Bendixson erasing rank is invariant under homeomorphism.

Thus $q(\mathcal{K}(G) \cap \text{Sub}_{N \leq}(G)) = q(\mathcal{K}(\text{Sub}_{N \leq}(G))) = \mathcal{K}(Q)$ and the restricted homeomorphism (3) is well-defined. The equality (4) also follows since, for $H \in \text{Sub}_{N \leq}(G)$,

$$\text{rk}_{\text{CB}}^{\times}(H; G) = \infty \Leftrightarrow H \in \mathcal{K}(G) \cap \text{Sub}_{N \leq}(G) \Leftrightarrow q(H) \in \mathcal{K}(Q) \Leftrightarrow \text{rk}_{\text{CB}}^{\times}(H/N; Q) = \infty$$

and, if $H \notin \mathcal{K}(G)$ then $H/N \notin \mathcal{K}(Q)$ so

$$\begin{aligned}
\mathrm{rk}_{\mathrm{CB}}^{\times}(H; G) &= \inf \left\{ \alpha \geq 1 : H \notin \mathrm{Sub}(G)^{(\alpha)} \right\} \\
&= \inf \left\{ \alpha \geq 1 : H \notin \mathrm{Sub}_{N \leq}(G)^{(\alpha)} \right\} \\
&= \inf \left\{ \alpha \geq 1 : q(H) \notin \mathrm{Sub}(Q)^{(\alpha)} \right\} \\
&= \mathrm{rk}_{\mathrm{CB}}^{\times}(H/N; Q).
\end{aligned}$$

If $\mathrm{Sub}(Q)$ is countable, then $\mathcal{K}(Q) = \emptyset$ and then $\mathrm{rk}_{\mathrm{CB}}^{\times}(Q) = \sup \{ \mathrm{rk}_{\mathrm{CB}}^{\times}(L; Q) : L \in \mathrm{Sub}(Q) \}$. This proves the inequality (5). \square

Propositions 3.3 and 3.5 imply that virtual algebraic fibering allows one to find finitely generated, infinite index subgroups which do not belong to the perfect kernel. More generally:

Corollary 3.6. *Let G be a countable group and $H \leq G$ be a finitely generated subgroup. Suppose that there exists a finite index subgroup $G' \leq G$ and a finitely generated subgroup $H' \leq H \cap G'$ such that there is a short exact sequence*

$$1 \longrightarrow H' \longrightarrow G' \longrightarrow Q \longrightarrow 1$$

where $\mathrm{Sub}(Q)$ is countable. Then $H \notin \mathcal{K}(G)$.

Proof. By Proposition 3.5, $\mathrm{Sub}_{H' \leq}(G')$ is a countable neighbourhood of H' . Let $P : G \rightarrow G'$ be defined by $P(L) = L \cap G'$. Then, by Proposition 3.3, $P^{-1}(\mathrm{Sub}_{H' \leq}(G'))$ is a countable neighbourhood of H . \square

3.2 Examples

Note that the existence of a short exact sequence with finitely generated kernel does not in general provide any information about subgroups which do not contain the kernel. In fact, there exist pairs $N \trianglelefteq G$ with N finite and normal in the countable group G such that $\{\mathrm{id}\} \in \mathcal{K}(G/N)$, yet the subgroup $\{\mathrm{id}\}$ is isolated in $\mathrm{Sub}(G)$.

Example 3.7 (Example with $\{\mathrm{id}\}$ isolated). In order to allow for the possibility that $\{\mathrm{id}\}$ be isolated, it is necessary that G contains no infinite order element. A very interesting example is a central extension by $\mathbb{Z}/p\mathbb{Z}$ of the Burnside group $B(m, p) := \langle h_1, h_2, \dots, h_m \mid g^p = 1, \forall g \rangle$. Let $m, p \in \mathbb{N}$, $m \geq 2$ and p a prime number ≥ 665 . There is a central extension

$$1 \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow G \longrightarrow B(m, p) \longrightarrow 1$$

in which all the elements have order p^2 except the elements of the center $\mathbb{Z}/p\mathbb{Z}$. Indeed, first take the Adian group $A(m, p)$ – this is a torsion free m -generated group whose center is an infinite cyclic group $\langle c \rangle$ and with quotient $A(m, p)/\langle c \rangle \simeq B(m, p)$ – and set $G = A(m, p)/\langle c^p \rangle$, see [Zel11, Example 1.25]. The p -th power of every element belongs to the center. It follows that every non-trivial subgroup contains the center. The trivial subgroup is thus isolated in $\mathrm{Sub}(G)$.

On the other hand, by a theorem of Širvanjan [Šir76], when $m \geq 2$ and p is larger than 665 and odd, $B(m, p)$ contains a copy of the countably infinite rank free Burnside group $B(\mathbb{N}, p) = \langle s_i, i \in \mathbb{N} \mid g^p = 1 \text{ for all } g \rangle$. Since $\{\mathrm{id}\} \in \mathcal{K}(B(\mathbb{N}, p)) = \mathrm{Sub}(B(\mathbb{N}, p))$, as shown in Proposition 3.8 below, then $\{\mathrm{id}\} \in \mathcal{K}(B(m, p))$. By Proposition 3.5, $\mathbb{Z}/p\mathbb{Z} \in \mathcal{K}(G)$.

Recall that the \mathfrak{R} -free group with generators indexed by I of a variety \mathfrak{R} of groups is the group $\mathfrak{F}_I^{\mathfrak{R}} = \langle s_i, i \in I \mid \mathfrak{R} \rangle$ where \mathfrak{R} represents a collection of relators of the form $R(g_1, g_2, \dots, g_p) = 1$ for all p -tuples (g_1, g_2, \dots, g_p) of elements of the (usual) free group on $(s_i)_{i \in I}$. Examples of \mathfrak{R} -free group $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$ of countably infinite rank include the free group \mathbf{F}_{∞} , the free abelian group $\oplus_{\mathbb{N}} \mathbb{Z}$, the free abelian group $\oplus_{\mathbb{N}} \mathbb{Z}/m\mathbb{Z}$ of exponent m , the free nilpotent group of class c , the free Burnside group $B(\mathbb{N}, p) = \langle s_i, i \in \mathbb{N} \mid g^p = 1, \text{ for all } g \rangle, \dots$ By *non-trivial* variety, we simply mean that $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$ is not finitely generated.

Proposition 3.8 (Free group of a variety). *Let $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}} = \langle s_1, s_2, \dots \mid \mathfrak{R} \rangle$ be the \mathfrak{R} -free group of countably infinite rank of a non-trivial variety \mathfrak{R} of groups. Then $\mathcal{K}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}) = \text{Sub}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}})$.*

Proof. For $I \subseteq \mathbb{N}$, consider the subgroup $H_I := \langle s_i : i \in I \rangle \leq \mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$, and let $\mathfrak{F}_I^{\mathfrak{R}} := \langle t_i : i \in I \mid \mathfrak{R} \rangle$ be the \mathfrak{R} -free group with generators indexed by I .

Consider the natural morphisms, well-defined by the universal property of the \mathfrak{R} -free groups, $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}} \xrightarrow{u} \mathfrak{F}_I^{\mathfrak{R}} \xrightarrow{v} \mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$ defined by $u(s_i) = t_i$ if $i \in I$ and $u(s_i) = \text{id}$ otherwise, and by $v(t_i) = s_i$.

Since v is the inverse of u when restricted to H_I , this ensures that $H_I \simeq \mathfrak{F}_I^{\mathfrak{R}}$ and $w_I := v \circ u$ is a *retraction* $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}} \rightarrow H_I$ (i.e. it is the identity on H_I). Similarly, when $I \subseteq J$, the obvious morphism $w_{I,J} : H_J \rightarrow H_I$ is a retraction.

Let L be a finitely generated subgroup of $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$. Then, there is a finite subset $I \subset \mathbb{N}$ such that $L \leq H_I$. For every $i \in \mathbb{N} \setminus I$, let $L_i = \langle L, s_i \rangle \leq H_{I \sqcup \{i\}}$. Observe, by non-triviality of the variety, that $s_i \notin H_I$ and thus $L_i \neq L$. We claim that the sequence $(L_i)_i$ tends to L when i tends to ∞ : If $g \in L_i$, then write it as a word m using elements of L and s_i . When $j > i$, if g continues to belong to L_j then under the retraction $H_{I \sqcup \{i,j\}} \rightarrow H_{I \sqcup \{j\}}$, we see that the word m continues to represent g even after the erasing of the letters s_i . Thus g can be written as a word using only elements of L , i.e. $g \in L$. It follows that L is a non-trivial limit of finitely generated subgroups of $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$ and thus all finitely generated subgroups belong to $\mathcal{K}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}})$. Now, if L is a non finitely generated subgroup of $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$, then it is a limit of finitely generated groups. We have proved that $\mathcal{K}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}) = \text{Sub}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}})$. \square

When $\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}}$ is abelian, its action on $\text{Sub}(\mathfrak{F}_{\mathbb{N}}^{\mathfrak{R}})$ is trivial, thus not topologically transitive, while $\mathbf{F}_{\infty} \curvearrowright \mathcal{K}(\mathbf{F}_{\infty})$ is highly topologically transitive (see Corollary 5.19).

Question 3.9 (Free Burnside groups and dynamics). For the free Burnside groups, $B(\mathbb{N}, p)$ of countably infinite rank and $B(m, p)$ of rank m :

1. Is the action $B(\mathbb{N}, p) \curvearrowright \text{Sub}(B(\mathbb{N}, p))$ topologically r -transitive for some r ?
2. For $m \geq 2$ and p such that $B(m, p)$ is infinite, what is perfect kernel $\mathcal{K}(B(m, p))$? How transitive is the corresponding action of $B(m, p)$?

3.3 Subgroups with Schreier graph quasi-isometric to \mathbb{Z}

We now consider subgroups which are not necessarily normal but whose ‘‘quotients’’ are nonetheless equipped with a strong geometric structure: namely those with Schreier graph quasi-isometric to \mathbb{Z} .

If G is a finitely generated group and H is a subgroup, we say that $H \backslash G$ is quasi-isometric to \mathbb{Z} if for a (equivalently any) Cayley graph X of G associated with a finite generating set, the **Schreier graph** $H \backslash X$ is quasi-isometric with \mathbb{Z} .

Theorem 3.10. *If $H \leq G$ be a pair of finitely generated groups such that $H \backslash G$ is quasi-isometric to \mathbb{Z} , then the Cantor-Bendixson erasing rank of H satisfies $\text{rk}_{\text{CB}}^{\times}(H; G) \leq 3$.*

Note that the above theorem does not hold if one assumes only that $H \backslash G$ is 2-ended. Indeed, if G is a hyperbolic surface group and $H \leq G$ is a cyclic subgroup then $H \backslash G$ is 2-ended but $H \in \mathcal{K}(G)$. It also does not hold if one removes the assumption the H is finitely generated: if G is a non-abelian free group and H is the kernel of a morphism onto \mathbb{Z} , then $H \in \mathcal{K}(G)$. We are however working on extending this result to finitely generated subgroups H such that $H \backslash G$ is quasi-isometric to \mathbb{Z}^d for $d \geq 2$.

Example 3.11. The trivial group $\{\text{id}\}$ in \mathbb{Z} and in the infinite dihedral group D_{∞} furnishes examples of Cantor-Bendixson erasing rank 2 and 3, respectively. This will be useful in Section 6.3.

Let G be a group with finite generating set S . Let $X := \text{Cay}(G, S)$ be the associated Cayley graph. The ball of radius R centered at g in $\text{Cay}(G, S)$ is denoted by $B_X(g, R)$.

For every tower of groups $H \leq K \leq G$, we shall consider the corresponding covering maps

$$X \xrightarrow{p_H} H \backslash X \xrightarrow{p_{K,H}} K \backslash X \xrightarrow{p_{G,K}} G \backslash X.$$

The idea of the proof is as follows. Given $H \leq G$ such that $H \backslash X$ is quasi-isometric to \mathbb{Z} , we will show that, for all but finitely many intermediate subgroups $H \leq K \leq G$, either K has finite index in G or the covering map $p_{K,H}$ folds the space $H \backslash X$ over, meaning that $K \backslash X$ is quasi-isometric to a half-line \mathbb{N} . In the latter case, we show that all but finitely many intermediate subgroups $K \leq L \leq G$ have finite index in G .

Proposition 3.12. *Let $H \leq G = \langle S \rangle$ be any subgroup. Let $x_0 \in H \backslash X$ and let $D > 0$ be a positive integer. There are at most $2^{|B_X(\text{id}, D)|}$ intermediate subgroups $H \leq K \leq G$ such that the fiber $p_{K,H}^{-1}(p_{K,H}(x_0))$ of $p_{K,H} : H \backslash X \rightarrow K \backslash X$ has diameter $\leq D$.*

Proof. Consider any such K . The finite set $p_{K,H}^{-1}(p_{K,H}(x_0)) = \{x_0, x_1, x_2, \dots, x_n\}$ is contained in the ball $B(x_0, D)$ of $H \backslash X$. Let $\tilde{x}_0 \in X$ be any pre-image of $x_0 \in H \backslash X$ under the covering map $p_H : X \rightarrow H \backslash X$. By lifting under p_H some geodesic paths from x_0 to x_i to geodesic paths from \tilde{x}_0 to \tilde{x}_i , one obtains a family of points $\{\tilde{x}_0, \tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n\}$ such that $p_H(\tilde{x}_i) = x_i$. Each \tilde{x}_i belongs to the ball $B_X(\tilde{x}_0, D)$ of X .

Let $g_0 = \text{id}, g_1, g_2, \dots, g_n \in G$ such that $g_i \tilde{x}_0 = \tilde{x}_i$. We claim that $K = \langle H, g_1, g_2, \dots, g_n \rangle$. In fact, $p_K(g_i \tilde{x}_0) = p_K(\tilde{x}_0)$ implies that $g_i \in K$. Thus $K \geq \langle H, g_1, g_2, \dots, g_n \rangle$. Since $p_K = p_{K,H} \circ p_H$, it follows that

$$K \cdot \tilde{x}_0 = H \cdot \{\tilde{x}_0, \tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n\} = \bigcup_{i=0}^n H \cdot g_i(\tilde{x}_0) = \left(\bigcup_{i=0}^n H g_i \right) \cdot \tilde{x}_0.$$

Since the action $G \curvearrowright X$ is free, it follows that $K \leq \langle H, g_1, g_2, \dots, g_n \rangle$.

Now let $A_D := \{\gamma_1, \gamma_2, \dots, \gamma_m\}$ be the set of elements of G such that $B_X(\tilde{x}_0, D) = \{\tilde{x}_0\} \cup \bigcup_{i=1}^m \gamma_i(\tilde{x}_0) \subseteq X$. If $K \leq G$ is a subgroup containing H such that the $p_{K,H}$ -fiber of $p_{K,H}(x_0)$ has diameter $\leq D$, then $K = \langle H, A_K \rangle$ for some subset $A_K \subseteq A_D$. The number of such subgroups K is thus bounded by the cardinality $2^{|B_X(\text{id}, D)|}$ of the power set of $B_X(\text{id}, D)$. \square

Lemma 3.13. *Consider subgroups $H \leq K \leq G$. Let $\bar{x} \in K \setminus X$ and $x_1, x_2 \in H \setminus X$ such that $p_{K,H}(x_1) = p_{K,H}(x_2) = \bar{x}$. Let $\bar{\delta}$ be a geodesic path in $K \setminus X$ starting at \bar{x} . Its pull-back δ_1 in $H \setminus X$ starting at x_1 is again a geodesic. If δ_1 meets the ball $B_{H \setminus X}(x_2, r)$ then $d(x_1, x_2) \leq 2r$.*

Proof. Let $z_2 \in \delta_1 \cap B_{H \setminus X}(x_2, r)$. Then, since (successively) $\delta_1, \bar{\delta}$ are geodesics, $p_{K,H}(x_1) = p_{K,H}(x_2)$ and $p_{K,H}$ does not increase distances, we have

$$d(x_1, z_2) = d(p_{K,H}(x_1), p_{K,H}(z_2)) = d(p_{K,H}(x_2), p_{K,H}(z_2)) \leq d(x_2, z_2) \leq r.$$

Thus, $d(x_1, x_2) \leq d(x_1, z_2) + d(z_2, x_2) \leq 2r$. \square

A connected graph (\mathcal{Y}, d) is C -quasi-isometric with \mathbb{Z} if there is a quasi-isometry $f : \mathbb{Z} \rightarrow \mathcal{Y}$, such that $\frac{1}{C}|i - j| - C \leq d(f(i), f(j)) \leq C|i - j| + C$ for every $i, j \in \mathbb{Z}$, and $d(z, f(\mathbb{Z})) \leq C$ for all $z \in \mathcal{Y}$. In this case, there is a constant C_1 (for instance $C_1 := 3C^3 + C^2 + 3C$) such that the complement of every ball of radius $\geq C_1$ has two infinite connected components.

Lemma 3.14. *Consider subgroups $H \leq K \leq G$. Assume $H \setminus X$ is C -quasi-isometric with \mathbb{Z} . Assume that $K \setminus X$ is infinite. Let $\bar{x} \in K \setminus X$ and $x_1, x_2 \in H \setminus X$ such that $p_{K,H}(x_1) = p_{K,H}(x_2) = \bar{x}$ and let $\bar{\delta}$ be an infinite geodesic path in $K \setminus X$ starting at \bar{x} . If the pull-backs δ_1, δ_2 starting at x_1, x_2 (respectively) point towards the same end ξ of $H \setminus X$, then*

$$d(\delta_1(t), \delta_2(t)) \leq C_2 = 2C_1$$

for every $t \in [0, \infty)$ where $\delta_1, \delta_2 : [0, \infty) \rightarrow H \setminus X$ are parametrised by arc length. In particular, $d(x_1, x_2) \leq C_2$.

Proof. Recall that the balls $B(x_i, C_1)$ both separate the space $H \setminus X$ and have two infinite components in their complement.

If $\delta_2 \cap B(x_1, C_1) = \emptyset$ then the end ξ is associated with the infinite connected component of $(H \setminus X) \setminus B(x_1, C_1)$ that contains δ_2 (and thus x_2 and $B(x_2, C_1)$). Then $B(x_2, C_1)$ separates $B(x_1, C_1)$ from ξ . Thus $\delta_1 \cap B(x_2, C_1) \neq \emptyset$. In conclusion, we have either $\delta_2 \cap B(x_1, C_1) \neq \emptyset$ or $\delta_1 \cap B(x_2, C_1) \neq \emptyset$. By Lemma 3.13, we conclude that $d(x_1, x_2) \leq 2C_1$. Then, $p_{K,H}(\delta_1(t)) = p_{K,H}(\delta_2(t)) = \bar{\delta}(t)$ are the initial point of infinite geodesic paths to which the above applies. \square

Proposition 3.15. *Consider subgroups $H \leq K \leq G$. Assume $H \setminus X$ is C -quasi-isometric with \mathbb{Z} . Then either $[K : H]$ is finite or $[G : K]$ is finite.*

Proof. Assume $[G : K]$ is infinite, i.e. $K \setminus X$ is infinite. Let $\bar{x} \in K \setminus X$ and let $\bar{\delta}$ be an infinite geodesic path in $K \setminus X$ starting at \bar{x} . If all the $p_{K,H}$ -pull-backs of $\bar{\delta}$ point towards the same end of $H \setminus X$, then $p_{K,H}^{-1}(\bar{x})$ has diameter bounded by C_2 (by Lemma 3.14). Assume now that there are two points $u_1, u_2 \in H \setminus X$ such that $p_{K,H}(u_i) = \bar{x}$ and such that the pull-backs of $\bar{\delta}$ starting at u_1, u_2 point towards different ends of $H \setminus X$. Then for all other points $u \in p_{K,H}^{-1}(\bar{x})$, the pull-back of $\bar{\delta}$ starting at u point towards the same end as either that starting at u_1 or that starting at u_2 . It follows that $d(u, \{u_1, u_2\}) \leq C_2$ (by Lemma 3.14). In any case, $p_{K,H}^{-1}(\bar{x})$ is finite. It follows that $[K : H]$ is finite. \square

Proposition 3.16. *Consider subgroups $H \leq K \leq G$. Assume $H \setminus X$ is C -quasi-isometric with \mathbb{Z} and that the index $[K : H]$ is finite, i.e. the fibers of $p_{K,H} : H \setminus X \rightarrow K \setminus X$ have cardinal*

$[K : H]$. Assume that a fiber of $p_{K,H}$ has diameter $> C_2 = 2C_1$. Then there are fibers of $p_{K,H}$ of arbitrarily large diameter and there are only finitely many subgroups $K \leq L \leq G$ such that $[L : K]$ is finite.

Proof. Let $\bar{x} \in K \setminus X$ and let $x_1, x_2 \in H \setminus X$ be two points such that $p_{K,H}(x_1) = p_{K,H}(x_2) = \bar{x}$ and $d(x_1, x_2) > C_2$. Observe that $K \setminus X$ is an infinite locally finite graph. Let $\bar{\delta}$ be a geodesic path in $K \setminus X$ starting at \bar{x} and let δ_1, δ_2 be its lifts in $H \setminus X$ starting at x_1 and x_2 respectively. By Lemma 3.14, δ_1, δ_2 point towards different ends ξ_1, ξ_2 , respectively, of $H \setminus X$. Then $p_{K,H}(\delta_1(t)) = p_{K,H}(\delta_2(t))$ for every $t \in [0, \infty)$ while the distance $d(\delta_1(t), \delta_2(t))$ tends to ∞ as t tends to ∞ : there is no bound on the diameter of the $p_{K,H}$ -fibers. Then, consider a subgroup L such that $K \leq L \leq G$ and $[L : K]$ is finite, thus $[G : L]$ is infinite. Consider the covering maps

$$H \setminus X \xrightarrow{p_{K,H}} K \setminus X \xrightarrow{p_{L,K}} L \setminus X,$$

and let us prove that there is a bound independent of L for the diameter of $p_{L,K}^{-1}(p_{L,K}(\bar{x}))$.

Let $z := p_{L,K}(\bar{x}) \in L \setminus X$ and let $B := (p_{L,K} \circ p_{K,H})^{-1}(z)$ be its pre-image under $p_{L,K} \circ p_{K,H}$. Let δ be a geodesic path in $L \setminus X$ starting at z and let δ_1, δ_2 be its lifts in $H \setminus X$ starting at x_1, x_2 respectively. Consider the partition $B = B_1 \sqcup B_2$ where B_i is the subset consisting of those b such that the lift of δ in $H \setminus X$ starting at b points towards ξ_i . Recall that $x_i \in B_i$. It follows from Lemma 3.14 that $B_i \subseteq B(x_i, C_2)$. Thus the diameter of B is bounded by $d(x_1, x_2) + 2C_2$. Since $p_{K,H}$ does not increase distances, it follows that $p_{L,K}^{-1}(p_{L,K}(\bar{x})) = p_{K,H}(B)$ has diameter bounded by $d(x_1, x_2) + 2C_2$. By Proposition 3.12, there are at most $2^{\lfloor B_X(\text{id}, d(x_1, x_2) + 2C_2) \rfloor}$ many such subgroups L . \square

Let us now give the proof of Theorem 3.10.

Proof of Theorem 3.10. Assume that $\text{rk}_{\text{CB}}^{\times}(H; G) > 2$. Let $(H_n)_n$ be a non-stationary sequence of subgroups of G that converges to H and such that $\text{rk}_{\text{CB}}^{\times}(H_n; G) \geq 2$. Since G is finitely generated, the finite index subgroups of G have erasing Cantor-Bendixson rank = 1, thus $[G : H_n] = \infty$. Since H is finitely generated, up to extraction one can assume that each H_n contains H , and so by Proposition 3.15, $[H_n : H] < \infty$. By Proposition 3.12, there are only finitely many subgroups $H \leq K \leq G$ for which the fibers of $p_{K,H}$ have diameter $\leq C_2 = 2C_1$. Thus up to extraction one can assume that each H_n gives rise to a $p_{H_n, H}$ -fiber of diameter $> C_2$.

Let us now approximate H_n . Let $(L_{n,j})_j$ be a non-stationary sequence tending to H_n when j tends to ∞ . Since $[H_n : H]$ is finite, H_n is finitely generated, thus we have $H \leq H_n \leq L_{n,j}$ for large enough j . By Proposition 3.16, there are only finitely many subgroups L such that $H \leq H_n \leq L \leq G$ and $[L : H_n] < \infty$, thus up to extraction, one can assume that $[L_{n,j} : H_n] = \infty$ for each j . Then, by Proposition 3.15 $[G : L_{n,j}] < \infty$. It follows that $\text{rk}_{\text{CB}}^{\times}(L_{n,j}; G) = 1$, so $\text{rk}_{\text{CB}}^{\times}(H_n; G) = 2$ for each n and $\text{rk}_{\text{CB}}^{\times}(H; G) = 3$. \square

4 Subgroup separable groups and the first ℓ^2 -Betti number

Let G be a countable group. A subgroup $H \leq G$ is **separable** if it is equal to the intersection of a sequence of finite index subgroups of G or equivalently if H is a closed subset of G with respect to the profinite topology. The group G is said to be **subgroup separable** if every

finitely generated subgroup is separable. Equivalently, G is subgroup separable if and only if the subspace $\text{Sub}_{[<\infty]}(G)$ of finite index subgroups of G is dense in $\text{Sub}(G)$, as observed in [GKM16]: the closure of the finite index subgroups contains all the finitely generated subgroups and these are dense in $\text{Sub}(G)$. Applied to the subgroup $\{\text{id}\}$ this property implies that G is residually finite. It is clear from the first definition that *subgroup separability passes to subgroups*. In addition it easily follows, for example via the profinite characterisation, that *subgroup separability is an invariant of commensurability*.

Subgroup separability was shown to hold for finitely generated free groups by M. Hall Jr in [Hal49] and for finitely generated orientable surface groups (and so all finitely generated Fuchsian groups) by P. Scott [Sco78].

Assuming G is a finitely generated subgroup separable group, the isolated points of $\text{Sub}(G)$ are exactly the finite index subgroups; i.e. H has finite index in G if and only if $\text{rk}_{\text{CB}}^\times(H; G) = 1$. We will show that $\text{Sub}(G)'$ has no isolated points in the case where G has positive first ℓ^2 -Betti number. We refer the reader to [CG86] for the precise definition of this invariant, but note that it verifies the following properties:

- $\beta_1^{(2)}(G) \leq \text{rk}(G)$ (this follows immediately from the definition).
- If $H \leq G$ is a finite index subgroup, then $\beta_1^{(2)}(H) = [G : H] \beta_1^{(2)}(G)$ (cf. [CG86, Proposition 2.6]).

The reader may observe that, in the proof of the following theorem, the first ℓ^2 -Betti number can be replaced by any group invariant $a(\cdot)$ such that, for any subgroup $H \leq G$, $a(H) \leq \text{rk}(H)$ and for any sequence $(H_n)_n$ of finite index subgroups of G , if $[G : H_n] \rightarrow \infty$ then $a(G) \rightarrow \infty$. In fact, the property we exploit here is that, for any sequence $(H_n)_n$ of finite index subgroups of G , if $[G : H_n] \rightarrow \infty$ then $\text{rk}(H_n) \rightarrow \infty$.

Theorem 4.1. *Let G be a subgroup separable group. If G is not finitely generated then $\text{Sub}(G)$ is a perfect space. If G is finitely generated with positive first ℓ^2 -Betti number then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ and thus $\text{rk}_{\text{CB}}(G) = 1$.*

Proof of Theorem 4.1. First suppose that G is not finitely generated and let $H \leq G$. If H is finitely generated then H has infinite index in G . Moreover, there is a sequence of finite index subgroups $(K_n)_n$ of G such that $H = \bigcap_n K_n$. In other words, the sequence of finite index subgroups $H_i = \bigcap_{n \leq i} K_n$ converges toward H and is non-stationary. Therefore H is not isolated in $\text{Sub}(G)$. If H is not finitely generated then it is a non-stationary limit of a sequence of finitely generated subgroups, so H is again not isolated. Thus $\text{Sub}(G)$ is perfect.

Now suppose that G is finitely generated and $\beta_1^{(2)}(G) > 0$. Every finite index subgroup of G is an isolated point of $\text{Sub}(G)$ so $\mathcal{K}(G) \subseteq \text{Sub}_{[\infty]}(G)$. Let $H \leq G$ be a subgroup with infinite index. If H is not finitely generated and $H = \langle s_n : n \in \mathbb{N} \rangle$ then $H = \lim_{n \rightarrow \infty} \langle s_1, \dots, s_n \rangle$ and, for each n , $\langle s_1, \dots, s_n \rangle \leq H$ has infinite index so H is not isolated in $\text{Sub}_{[\infty]}(G)$. Next, suppose that H has a finite generating set of cardinality r . By subgroup separability of G , the subgroup H is the intersection of countably many finite index subgroups of G and, since finite intersections of finite index subgroups have finite index, H is the limit of a strictly decreasing sequence $(K_n)_{n \in \mathbb{N}}$ of finite index subgroups of G . For each n , let $H_n := \langle H, k_n \rangle$ for some $k_n \in K_n \setminus H$. Since $H \supsetneq H_n \leq K_n$ for all n it follows that $H_n \rightarrow H$ non trivially. It remains to show that H_n has infinite index in G for almost all n .

Since $(K_n)_{n \in \mathbb{N}}$ is a non-stationary nested sequence of finite index subgroups, the sequence $([G : K_n])_{n \in \mathbb{N}}$ of indices must tend towards infinity. In particular, $\beta_1^{(2)}(G)[G : K_n] > r + 1$, for almost all n . For all such n , if H_n has finite index in G then

$$\beta_1^{(2)}(H_n) = \beta_1^{(2)}(G)[G : H_n] \geq \beta_1^{(2)}(G)[G : K_n] > r + 1.$$

But H_n has a generating set of cardinality $r + 1$ so $r + 1 \geq \beta_1^{(2)}(H_n)$ and this is a contradiction. Hence, for almost all n , H_n has infinite index as required. It follows that $\text{Sub}_{[\infty]}(G)$ has no isolated points. \square

Remark 4.2. Kielak showed in [Kie20] that if G is a finitely generated group which is virtually residually finite rationally solvable (or RFRS) in the sense of [Ago08] then $\beta_1^{(2)}(G) = 0$ if and only if G virtually fibers (i.e. G has a finite index subgroup which surjects onto \mathbb{Z} with finitely generated kernel). This allows us to sharpen our result in the following sense:

If G is finitely generated, subgroup separable and virtually RFRS then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ if and only if $\beta_1^{(2)}(G) > 0$.

One direction is given by Theorem 4.1 while the other follows from Corollary 3.6 since it implies that the kernel of a virtual fibering is an infinite index subgroup which is not contained in the perfect kernel of G . In particular, we characterise the property of having maximal perfect kernel in terms of the first ℓ^2 -Betti number for a large class of subgroup separable groups, including all those which virtually embed into right-angled Artin groups.

The following corollary is a generalisation of a well-known fact about free groups (see e.g. [CGLM20, Proposition 2.1]).

Corollary 4.3. *If G is a virtually free group of rank ≥ 2 then*

$$\mathcal{K}(G) = \begin{cases} \text{Sub}_{[\infty]}(G) & \text{if } G \text{ is finitely generated} \\ \text{Sub}(G) & \text{otherwise.} \end{cases}$$

Proof. If \mathbf{F}_r is a free group of rank $r \geq 2$ then $\beta_1^{(2)}(\mathbf{F}_r) = r - 1 > 0$. Moreover it was shown in [Hal49] that \mathbf{F}_r is subgroup separable. This implies that any group G containing \mathbf{F}_r as a finite index subgroup is subgroup separable and has positive first ℓ^2 -Betti number so the result follows by Theorem 4.1. \square

Corollary 4.4. *If G is virtually a finitely generated non-abelian limit group, then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$.*

Proof. Limit groups are subgroup separable by [Wil08] and a finitely generated, non-abelian limit group has positive first ℓ^2 -Betti number by [Pic06, p. 645, l. 2-3]. The result therefore follows from Theorem 4.1. \square

In the special case of surface groups, the perfect kernel was also described using a different proof in unpublished work by A. Carderi, the second author and F. Le Maître.

Example 4.5 (Product of free groups over a cyclic subgroup). Let $G = \mathbf{F}_p *_Z \mathbf{F}_q$ for some finite $p, q \geq 2$. By [BBS84], free products of subgroup separable groups over cyclic subgroups are subgroup separable. Moreover, using the Mayer-Vietoris formula from [CG86, p. 204, §4], the first ℓ^2 -Betti number of G satisfies $\beta_1^{(2)}(G) = p + q - 2 > 0$ so, by Theorem 4.1, $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$. For information about the dynamics, see Corollary 5.23.

5 Actions on trees

We will refer throughout to the geometric realisation of an abstract graph as a graph. More precisely, a **graph** is a one-dimensional cell complex X . We denote by $V(X)$ its set of vertices (or 0-cells) and by $E(X)$ its set of edges (or 1-cells). For each $e \in E(X)$ let $\varphi_e : \partial e \rightarrow V(X)$ be the corresponding attaching map and define maps:

$$o, t : E(X) \rightarrow V(X)$$

where for each $e = [0, 1] \in E(X)$ its **endpoints** are the vertices $o(e) = \varphi_e(0)$ and $t(e) = \varphi_e(1)$. Note that X is endowed with a natural orientation where each edge e is directed from $o(e)$ to $t(e)$. A **path** P in X is a sequence of edges $(e_n)_{n \in N}$, where $N = \mathbb{N} = \mathbb{N}_{>0}$ or $N = \{1, \dots, k\}$ for some $k \in \mathbb{N}$, such that $e_n = \gamma([n-1, n])$ for each $n \in N$ and for some continuous map $\gamma : \cup_{n \in N} [n-1, n] \rightarrow X$. If N has cardinality $k \in \mathbb{N}$ then k is the **length** of P and if $N = \mathbb{N}$ we say that P is **infinite**. If γ is injective then we say that P is **simple**.

If X, Y are graphs and $\varphi : X \rightarrow Y$ is a bijection then φ is an isometry if it preserves the cellular structure of X . It follows that φ preserves the length metrics d_X and d_Y on X and Y respectively. All actions on graphs are assumed to be by orientation preserving isometries. A **tree** is a simply connected graph.

Recall that the action of a group G on a tree Y is **minimal** if Y does not contain any proper G -invariant subtree. The action is **reducible** if one of the following holds: (1) there is a global fixed point; (2) G fixes an end of Y ; or (3) there is a G -invariant line in Y . Otherwise, the action is **irreducible** (aka of general type). If x is a vertex or an edge of Y , we denote its stabiliser under the action of a subgroup $H \leq G$ by $\text{Stab}_H(x)$.

The structure theorem of Bass-Serre Theory [Ser80, §5.4, Theorem 13] implies that a group G admits a minimal action on a tree with a single orbit of edges if and only if it splits non-trivially as an amalgamated free product $G = A *_C B$ or a HNN extension $G = A *_C$. The G -tree Y associated with such a splitting is called its Bass-Serre tree. In the former case G acts on Y with two orbits of vertices, C is the stabiliser of an edge and A, B are the stabilisers of its endpoints and in the latter G acts with a single orbit of vertices, C is the stabiliser of an edge and A is the stabiliser of one of its endpoints.

We will make use of the following standard characterisation of irreducibility.

Proposition 5.1. *Consider a minimal action $G \curvearrowright T$ on a tree that is transitive on the set of edges. The action is irreducible if and only if, for some (equivalently every) edge e of T with endpoints v and w , the indices of the stabilisers satisfy (up to relabelling v and w):*

1. $[\text{Stab}_G(v) : \text{Stab}_G(e)] \geq 2$ and $[\text{Stab}_G(w) : \text{Stab}_G(e)] \geq 2$, when $G \backslash T$ has one vertex.
2. $[\text{Stab}_G(v) : \text{Stab}_G(e)] \geq 3$ and $[\text{Stab}_G(w) : \text{Stab}_G(e)] \geq 2$, when $G \backslash T$ has two vertices.

Proof. Recall that for an edge e of T and an endpoint v of e , the set of edges in the G -orbit of e and adjacent to v is indexed by the cosets $\text{Stab}_G(v)/\text{Stab}_G(e)$.

Assume $G \backslash T$ has one edge and two vertices. If $[\text{Stab}_G(v) : \text{Stab}_G(e)] = [\text{Stab}_G(w) : \text{Stab}_G(e)] = 2$, then T is a line and the action is reducible. If $[\text{Stab}_G(v) : \text{Stab}_G(e)] = 1$, then v has degree 1 in T : this contradicts the minimality of the action.

Assume $G \backslash T$ has one edge and one vertex. If $[\text{Stab}_G(v) : \text{Stab}_G(e)] = 1$, then up to reversing the orientation of all the edges, every vertex of T is the startpoint of a single edge:

for every $x \in V(T)$, there is a single edge $f \in E(T)$ such that $o(f) = x$. There is a unique infinite simple path $P = (e_n)_{n \in \mathbb{N}}$ with startpoint x that follows the orientations of the edges. For any two vertices, the corresponding infinite simple positive path must eventually agree. They thus define the same end of T . This uniquely defined end is fixed by G and the action on T is reducible.

We have proved that irreducibility of the action implies the desired conditions on the stabilisers.

Conversely, if T admits an edge e with an endpoint v such that the index of the stabiliser satisfies $[\text{Stab}_G(v) : \text{Stab}_G(e)] \geq 3$, it follows that T is not a line: v has degree ≥ 3 . If $G \curvearrowright T$ has only one vertex and $[\text{Stab}_G(v) : \text{Stab}_G(e)] = [\text{Stab}_G(w) : \text{Stab}_G(e)] = 2$, then every vertex has degree ≥ 4 , thus T is not a line. There is no global fixed point or G -invariant line in T by minimality of the action. Moreover, if $P = (e_n)_{n \in \mathbb{N}}$ is an infinite simple path starting at v_0 and $g \in \text{Stab}_G(v_0) \setminus \text{Stab}_G(e_1)$ then gP is an infinite path in T whose intersection with P is $\{v_0\}$. Therefore P and gP define distinct ends of T so no end of T is fixed by G . \square

Let T be a tree and $\varphi : T \rightarrow T$ be a graph automorphism. Recall that φ is either **elliptic** (i.e. φ fixes a point in T) or **hyperbolic** (i.e. the translation length of φ is realised in T and is strictly positive) [Ser80, Proposition 2.4]. In the later case, there is a unique bi-infinite line $\text{Axis}(\varphi)$ in T on which φ acts by translation.

Lemma 5.2. *Let $G \curvearrowright T$ be a minimal irreducible action of a countable group. Let e be an edge of T and Z be one of the two connected components of $T \setminus \{e\}$. Let L be a finite segment of edges $[f_1, f_2]$. Then there are three hyperbolic elements $\gamma_0, \gamma_1, \gamma_2 \in G$ such that*

- $\text{Axis}(\gamma_0)$ contains e ;
- $\text{Axis}(\gamma_1)$ is contained in Z ;
- $\text{Axis}(\gamma_0) \cap \text{Axis}(\gamma_1)$ is non-empty and bounded (the hyperbolic elements are “transverse”);
- the segment $\gamma_2(L)$ is contained in Z and $\gamma_2 \cdot f_1$ separates $\gamma_2 \cdot f_2$ from e .

Proof. By shrinking to a point each connected component of $T \setminus G \cdot e$, the complement in T of the G -orbit of e , we obtain a tree T' equipped with a G -action that is transitive on the set of edges and for which the map $T \rightarrow T'$ is G -equivariant. By minimality of the action $G \curvearrowright T$, the convex hull of $G \cdot e$ is the whole of T ; thus $G \curvearrowright T'$ is minimal and irreducible and both actions have the same kernel. If γ_0, γ_1 satisfy the conditions of the lemma for T' , then their action on T also satisfies these conditions.

Thus, one can start by assuming that T has a single orbit of edges. There is a hyperbolic element γ_0 whose axis contains e : by irreducibility $G \curvearrowright T$ admits some hyperbolic element h_0 , and for any edge $f \in \text{Axis}(h_0)$, there is $h_1 \in G$ such that $h_1 \cdot f = e$; then $h_1(\text{Axis}(h_0))$ is the axis of $\gamma_0 := h_1 h_0 h_1^{-1}$ and it contains e .

Recall from [Ser80, I.6.4, p. 62] that two edges f and f' are **coherent** if they are oriented in the same way in the unique geodesic segment which connects them. Let a, b be edges of $Z \setminus \text{Axis}(\gamma_0)$ that are adjacent to $\text{Axis}(\gamma_0)$ but at different vertices of $\text{Axis}(\gamma_0)$. If a and b are not coherent, one replaces b by an edge b' that is adjacent to b , that is separated from $\text{Axis}(\gamma_0)$ by b and that is coherent with a (every vertex is adjacent to at least two incoherent edges – see Proposition 5.1). Thus, one can take a and b to be coherent. Choose an element $\gamma_1 \in G$ sending a to b : then by [Ser80, Corollary, p. 63] γ_1 is hyperbolic and $[a, b]$ belongs to its axis. In particular $\text{Axis}(\gamma_1)$ has a non-trivial and bounded intersection with $\text{Axis}(\gamma_0)$. Moreover $[a, b]$ separates $\text{Axis}(\gamma_1) \setminus [a, b]$ from e , in particular, $\text{Axis}(\gamma_1) \subset Z$.

There is an integer r such that $\gamma_0^r(L) \subset Z$. If $\gamma_0^r \cdot f_1$ separates $\gamma_0^r \cdot f_2$ from e , set $\gamma_2 := \gamma_0^r$. Otherwise, the connected component Z_2 of $T \setminus \{\gamma_0^r \cdot f_1\}$ that does not contain $\gamma_0^r \cdot f_2$ is entirely contained in Z . Choose $\gamma_1' \in G$ whose axis is entirely contained in Z_2 . Then the path from $\gamma_1' \gamma_0^r \cdot f_2$ to e traverses $\gamma_1' \gamma_0^r \cdot f_1$ and $\gamma_2 := \gamma_1' \gamma_0^r$ satisfies the condition. It is easy to check that γ_2 is hyperbolic (the projection of L onto $\text{Axis}(\gamma_0)$ belongs to the axis of γ_2). \square

The **kernel** of an action $G \curvearrowright T$ is the normal subgroup consisting of the elements which fix every vertex of T .

Proposition 5.3. *If $G \curvearrowright T$ is a minimal irreducible action on a tree with finite kernel N , then $\text{Sub}(N) \subseteq \mathcal{K}(G)$.*

Proof. By [CM87, Theorem 2.7] the irreducibility of the action implies that G has a free subgroup \mathbf{F}_2 of rank 2. Therefore $\text{Sub}(N) \subseteq \mathcal{K}(G)$ by Proposition 3.4. \square

5.1 Main technical dendrological lemma

Lemma 5.4 (Main technical dendrological lemma). *Let $G \curvearrowright T$ be a minimal action on a tree T . Let $\Lambda \leq G$ be a subgroup. Assume that Λ admits a proper invariant subtree $T_\Lambda \subsetneq T$ for which there are sequences $(H_n)_n$, $(S_n)_n$ and $(e_n)_n$ such that for every n*

1. H_n is a subgroup of G ;
2. S_n is an H_n -invariant subtree of T ;
3. $e_n \in E(T)$ is an edge such that:
 - (a) e_n is at distance greater than $n + 1$ from T_Λ
 - (b) e_n separates T_Λ from S_n ;
 - (c) $\text{Stab}_\Lambda(e_n) = \text{Stab}_{H_n}(e_n)$, thus $\Lambda \cap H_n = \text{Stab}_\Lambda(e_n)$;
 - (d) $\text{Stab}_{H_n}(e_n) \neq H_n$;
 - (e) The Λ -orbit of e_n does not separate T_Λ from the other e_k , $k \in \mathbb{N} \setminus \{n\}$.

Let $\Lambda_n := \langle \Lambda, H_n \rangle$. Let \mathcal{L}_T be the set of subgroups of G that satisfy assumptions 1-3. Then:

- (I) Each Λ_n splits as $\Lambda_n = \Lambda *_{\text{Stab}_\Lambda(e_n)} H_n$ and $\text{Stab}_{\Lambda_n}(e_n) = \text{Stab}_\Lambda(e_n)$.
More generally, $\text{Stab}_{\Lambda_n}(e_k) = \text{Stab}_\Lambda(e_k)$ for all $k \in \mathbb{N}$ and $n \in \mathbb{N}$.
- (II) The sequence $(\Lambda_n)_n$ tends non trivially to Λ .
- (III) Each Λ_n belongs to \mathcal{L}_T , witnessed by a subsequence of $(H_n)_n$, $(S_n)_n$ and $(e_n)_n$.
- (IV) The set \mathcal{L}_T has no isolated points for its induced topology. In particular $\mathcal{L}_T \subseteq \mathcal{K}(G)$.
- (V) For every $\Lambda \in \mathcal{L}_T$, the quotient $\Lambda \backslash T$ is infinite.

Proof. Step (I). We fixed an integer n . We will first construct a tree A_n on which the subgroup $\Lambda_n := \langle \Lambda, H_n \rangle$ of G acts. Let v_n be the endpoint of e_n nearest to T_Λ and let w_n be the other endpoint of e_n .

Denote by $T_{n,\Lambda}^*$ the connected component of $T \setminus \{\Lambda \cdot e_n\}$ that contains T_Λ and by $T_{H_n}^*$ the connected component of $T \setminus \{H_n \cdot e_n\}$ that contains S_n .

Notice (by Assumption (3b)) that $T_{n,\Lambda}^*$ and $T_{H_n}^*$ are disjoint subtrees of T that are Λ -invariant and H_n -invariant respectively. Moreover, e_n connects them. Also observe for later use that by Assumption (3e), all the e_k , $k \in \mathbb{N} \setminus \{n\}$ are contained in $T_{n,\Lambda}^*$.

By an immediate induction based on the length of an element of $\Lambda_n = \langle \Lambda, H_n \rangle$ as a product of elements from Λ and H_n , we obtain that the Λ_n -orbit of e_n splits T into connected components that are either a Λ_n -image of $T_{n,\Lambda}^*$ or a Λ_n -image of $T_{H_n}^*$. Contracting each of these connected components to a point, we obtain a Λ_n -equivariant map

$$P_n : T \rightarrow A_n$$

from T to a tree A_n equipped with the induced Λ_n -action, with a single Λ_n -orbit of edges and two Λ_n -orbits of vertices.

Let \bar{v}_n be the vertex corresponding to the connected component $T_{n,\Lambda}^*$ and \bar{w}_n be the vertex corresponding to the connected component $T_{H_n}^*$. Let \bar{e}_n be the image of e_n ; its endpoints are \bar{v}_n and \bar{w}_n . It follows from general Bass-Serre theory that Λ_n splits as

$$\Lambda_n = \text{Stab}_{\Lambda_n}(\bar{v}_n) *_{\text{Stab}_{\Lambda_n}(\bar{e}_n)} \text{Stab}_{\Lambda_n}(\bar{w}_n). \quad (6)$$

Our goal is now to identify this splitting with the splitting $\Lambda_n = \Lambda *_{\text{Stab}_\Lambda(e_n)} H_n$ of (I).

The projection map $P_n : T \rightarrow A_n$ is injective on the Λ_n -orbit of e_n , thus the stabilisers of e_n and \bar{e}_n are the same for Λ and H_n , respectively:

$$\text{Stab}_\Lambda(\bar{e}_n) = \text{Stab}_\Lambda(e_n) \quad \text{and} \quad \text{Stab}_{H_n}(\bar{e}_n) = \text{Stab}_{H_n}(e_n).$$

Thus, by Assumption (3c), $\text{Stab}_\Lambda(\bar{e}_n) = \text{Stab}_{H_n}(\bar{e}_n)$. By definition $\text{Stab}_{\Lambda_n}(\bar{e}_n) \cap \Lambda = \text{Stab}_\Lambda(\bar{e}_n)$ and $\text{Stab}_{\Lambda_n}(\bar{e}_n) \cap H_n = \text{Stab}_{H_n}(\bar{e}_n)$. It follows that:

$$\text{Stab}_{\Lambda_n}(\bar{e}_n) \cap \Lambda = \text{Stab}_{\Lambda_n}(\bar{e}_n) \cap H_n = \Lambda \cap H_n. \quad (7)$$

Since $T_{n,\Lambda}^*$ is Λ -invariant and $T_{H_n}^*$ is H_n -invariant in T , the stabilisers of their images $P(T_{n,\Lambda}^*) = \bar{v}_n$ and $P(T_{H_n}^*) = \bar{w}_n$ for the Λ_n -action on A_n satisfy

$$\Lambda \leq \text{Stab}_{\Lambda_n}(\bar{v}_n) \quad \text{and} \quad H_n \leq \text{Stab}_{\Lambda_n}(\bar{w}_n). \quad (8)$$

We shall make use of the normal form theorem [LS77, Theorem 2.6, Section IV.2, p. 187] for the amalgamated free product splitting (6). Since $\langle \Lambda, H_n \rangle = \Lambda_n$, every non-trivial element $g \in \text{Stab}_{\Lambda_n}(\bar{v}_n)$ can be written in the form $g = c_1 c_2 \cdots c_j$ where

- each c_i is in one of the “factor” subgroups Λ or H_n ,
- successive c_i, c_{i+1} come from different factors, and
- if $j > 1$ then no c_i is in the intersection $\Lambda \cap H_n$.

Then (by (8) and (7)) $c_1 c_2 \cdots c_j$ is a reduced sequence for the splitting (6). On the other hand, $g^{-1} c_1 c_2 \cdots c_j = 1$ is not reduced for the splitting (6). If $j > 1$, then $g^{-1} c_1 \in \text{Stab}_{\Lambda_n}(\bar{e}_n)$, and successively we get $g^{-1} c_1 c_2 \cdots c_{j-1} \in \text{Stab}_{\Lambda_n}(\bar{e}_n)$. In the end, c_j also belongs to $\text{Stab}_{\Lambda_n}(\bar{e}_n)$,

and by (7), $c_j \in \Lambda \cap H_n$, a contradiction. Thus $j = 1$ and $g = c_1 \in \Lambda$ and $\Lambda = \text{Stab}_{\Lambda_n}(\bar{v}_n)$. Similarly, the symmetric argument with the other factor shows that $H_n = \text{Stab}_{\Lambda_n}(\bar{w}_n)$. We have proved that

$$\text{Stab}_{\Lambda_n}(\bar{v}_n) = \Lambda, \quad \text{Stab}_{\Lambda_n}(\bar{w}_n) = H_n \quad \text{and} \quad \text{Stab}_{\Lambda_n}(e_n) = \text{Stab}_{\Lambda}(e_n). \quad (9)$$

and the splitting (6) becomes

$$\Lambda_n = \Lambda *_{\text{Stab}_{\Lambda}(e_n)} H_n. \quad (10)$$

The ‘‘More generally’’ part of (I), i.e. $\text{Stab}_{\Lambda_n}(e_k) = \text{Stab}_{\Lambda}(e_k)$ for all the other $k \in \mathbb{N} \setminus \{n\}$, is proved later as Equation (11).

Step (II). All the Λ_n contain Λ , thus in order to show that $(\Lambda_n)_n$ tends to Λ , it is enough to prove that for every $g \in G$, there is n_0 such that if $g \in \Lambda_n$ for some $n \geq n_0$, then $g \in \Lambda$.

Let $g \in G$, let v be a vertex v in T_{Λ} and let n_0 be the distance in T between v and $g \cdot v$. Assume that $g \in \Lambda_n$ for some $n \geq n_0$. By Assumption (3a) the n -neighbourhood of T_{Λ} is contained in $T_{n,\Lambda}^*$, thus both v and $g \cdot v$ belong to $T_{n,\Lambda}^*$. It follows that their images in A_n both equal \bar{v}_n . By equivariance of the quotient map $T \rightarrow A_n$, the element g fixes the vertex \bar{v}_n . Since $\text{Stab}_{\Lambda_n}(\bar{v}_n) = \Lambda$ (by Equation (9)), it follows that $g \in \Lambda$. We have proved that $\Lambda_n \rightarrow \Lambda$ as $n \rightarrow \infty$.

By Assumption (3d), the injection $\Lambda \leq \Lambda_n = \Lambda *_{\text{Stab}_{\Lambda}(e_n)} H_n$ is strict, so the convergence of the sequence $(\Lambda_n)_n$ is non-trivial.

Step (III). Let us show that $\Lambda_n \in \mathcal{L}_T$.

The invariant tree we propose for T_{Λ_n} is the Λ_n -saturation of the convex hull of $T_{\Lambda} \cup S_n$:

$$T_{\Lambda_n} := \Lambda_n \cdot \text{Hull}(T_{\Lambda} \cup S_n).$$

As for the sequences of subgroups $(H'_k)_k$, subtrees $(S'_k)_k$, and edges $(e'_k)_k$ associated to Λ_n , we will take subsequences of those associated with Λ , obtained by shifting the indices by r_n , the distance between T_{Λ} and v_n , i.e. $r_n := d(T_{\Lambda}, v_n)$, $H'_k := H_{k+r_n}$, $S'_k := S_{k+r_n}$ and $e'_k := e_{k+r_n}$.

As observed earlier, all the e_k , for $k \in \mathbb{N} \setminus \{n\}$ are contained in $T_{n,\Lambda}^*$ by Assumption (3e). If an element g of Λ_n were to send some e_i to $g \cdot e_i$ so that $g \cdot e_i$ separated some e_j from T_{Λ_n} (for $i, j \neq n$), then $g \cdot e_i$ would also belong to $T_{n,\Lambda}^*$ and g would fix the vertex \bar{v}_n of A_n . Thus g would belong to Λ (by Equation (9)). But Assumption (3e) for Λ and all n prevents the Λ -orbit of e_i from separating T_{Λ} and e_j . Thus the sequence $(e'_k)_k$ satisfies Assumption (3e) for Λ_n , and in particular the Λ_n -invariant tree T_{Λ_n} is strictly contained in T . It also follows that e'_k separates T_{Λ_n} from S'_k , i.e. Assumption (3b) for Λ_n . Since $T_{\Lambda_n} \cap T_{n,\Lambda}^*$ is contained in the r_n -neighbourhood of T_{Λ} , this shift in the indices permits to check Assumption (3a) for Λ_n .

An element g of Λ_n that fixes some e_k , for $k \in \mathbb{N} \setminus \{n\}$ (which again is an edge of $T_{n,\Lambda}^*$) belongs to the stabiliser of \bar{v}_n , thus to Λ by Equation (9). Thus for all $k \in \mathbb{N}$:

$$\text{Stab}_{\Lambda_n}(e_k) = \text{Stab}_{\Lambda}(e_k). \quad (11)$$

Assumption (3c) for Λ ensures that $\text{Stab}_{\Lambda}(e_k) = \text{Stab}_{H_k}(e_k)$, so $\text{Stab}_{\Lambda_n}(e_k) = \text{Stab}_{H_k}(e_k)$. Thus the sequence $(e'_k)_k$ satisfies Assumption (3c) for Λ_n . Assumption (3d) for Λ_n , i.e.

$\text{Stab}_{H'_k}(e'_k) \neq H'_k$ is immediate from the analogous condition $\text{Stab}_{H_{k+r_n}}(e_{k+r_n}) \neq H_{k+r_n}$ for Λ . We have proved that $\Lambda_n \in \mathcal{L}_T$.

Therefore, every point of \mathcal{L}_T is a non-trivial limit of points of \mathcal{L}_T . The proof of Item (IV) follows immediately.

For $\Lambda \in \mathcal{L}_T$ and any edge $e \in E(T)$, the distance from e to T_Λ is an invariant of its Λ -orbit $\Lambda \cdot e$. This is precisely the distance in $\Lambda \backslash T$ between $\Lambda \backslash T_\Lambda$ and the image of e in $\Lambda \backslash T$. By Assumption (3a), the sequence of edges (e_n) thus delivers infinitely many edges in $\Lambda \backslash T$. This proves Item (V), and concludes the proof of the main technical dendrological Lemma 5.4. \square

5.2 Main dendrological theorems

Theorem 5.5. *Let $G \curvearrowright T$ be a minimal and irreducible action with finite kernel N on a tree T . Let $\Lambda \leq G$ be a subgroup and suppose that there exists an edge $e_\Lambda \in E(T)$ such that $\text{Stab}_\Lambda(e_\Lambda) = \Lambda \cap N$ and one of the connected component of $T \setminus \{e_\Lambda\}$ contains a Λ -invariant subtree $T_\Lambda \subseteq T$. Then Λ belongs to $\mathcal{K}(G)$.*

More precisely, the set \mathcal{N}_T of subgroups that satisfy the assumptions of this theorem for the action $G \curvearrowright T$ has no isolated points with respect to its induced topology.

Proof. We will show that Λ satisfies the conditions of Lemma 5.4.

Let w_0 be the endpoint of e_Λ that is contained in the component Z of $T \setminus e_\Lambda$ which does not contain T_Λ , and let w_1 be the other endpoint of e_Λ . Since e_Λ separates Z from a Λ -invariant subtree, it follows that if u_1, u_2 are vertices of Z and $\lambda \in \Lambda$ such that $\lambda \cdot u_1 = u_2$, then λ fixes w_0 and e_Λ , so $\lambda \in \Lambda \cap N$. In particular λ belongs to the kernel of the action, thus the quotient map $T \rightarrow \Lambda \backslash T$ is injective on Z . Moreover, (taking $u_1 = u_2$) for every vertex $v \in V(Z)$ and every edge $e \in E(Z)$ the stabilisers satisfy:

$$\text{Stab}_\Lambda(v) = \Lambda \cap N = \text{Stab}_\Lambda(e). \quad (12)$$

By Lemma 5.2, there is a hyperbolic element g whose axis is contained in Z . Thus the sequence of edges $(e_n)_{n \in \mathbb{N}} := (g^n \cdot e_\Lambda)_{n \in \mathbb{N}}$, satisfies two conditions of Lemma 5.4: (3a) (distance to T_Λ) and (3e) (the Λ -orbit of the edge e_n intersects Z only in e_n , thus does not separate any other e_k from T_Λ).

By Lemma 5.2, there is a hyperbolic element h_n for each $n \in \mathbb{N}$ whose axis is contained in the connected component of $T \setminus \{e_n\}$ that does not contain T_Λ . We define the subtree S_n as the axis of h_n . The sequence $(e_n)_n$ satisfies (3b) of Lemma 5.4.

Since N is finite and normal in G , there is a power r_n of h_n which commutes with N . Let H_n be the subgroup generated by $h_n^{r_n}$ and $\Lambda \cap N$:

$$H_n := \langle h_n^{r_n}, \Lambda \cap N \rangle.$$

Since $h_n^{r_n}$ is hyperbolic and commutes with $\Lambda \cap N$ and since $\Lambda \cap N$ is in the kernel of the action, $g \in H_n$ fixes e_n (or any other edge) if and only if $g \in \Lambda \cap N$. Thus $\text{Stab}_{H_n}(e_n) = \Lambda \cap N = \text{Stab}_\Lambda(e_n)$. This shows condition (3c) of Lemma 5.4.

As for condition (3d), it is clear since $\text{Stab}_{H_n}(e_n)$ is contained in the kernel of the action $G \curvearrowright T$ while H_n contains the hyperbolic element $h_n^{r_n}$. Thus Λ satisfies the conditions of Lemma 5.4.

By Conclusion (I) of Lemma 5.4, the group Λ is the non-trivial limit of

$$\Lambda_n := \langle \Lambda, H_n \rangle = \Lambda *_{(\Lambda \cap N)} H_n.$$

Since $\Lambda_n \cap N$ is contained in the edge stabiliser $\Lambda \cap N$ which is in turn contained in N , we have $\Lambda \cap N = \Lambda_n \cap N$. By Conclusion (III) of Lemma 5.4, Λ_n belongs to \mathcal{L}_T for a subsequence of the $(e_n)_n$ and a proper subtree $T_{\Lambda_n} \subsetneq T$ that does not contain these edges. For any $k \in \mathbb{N}$, we have $\text{Stab}_{\Lambda_n}(e_k) = \text{Stab}_{\Lambda}(e_k)$ (by the ‘‘More generally’’ part of Conclusion (I) of Lemma 5.4) and, since all the e_k belong to $E(Z)$, Equation (12) gives $\text{Stab}_{\Lambda_n}(e_k) = \Lambda \cap N = \Lambda_n \cap N$. Thus each Λ_n satisfies the assumptions of Theorem 5.5, thereby completing the proof that \mathcal{N}_T has no isolated points. This concludes the proof of Theorem 5.5. \square

If $G \curvearrowright T$ is a minimal and irreducible action on a tree, let us denote by

$$\text{Sub}_{|\bullet \setminus T|_\infty}(G) := \{\Lambda \in \text{Sub}(G) : |\Lambda \setminus T| = \infty\}$$

the set of subgroups whose action on T has infinitely many orbits of edges. Observe that $\text{Sub}_{|\bullet \setminus T|_\infty}(G)$ as well as its closure $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$ are invariant under the action $G \curvearrowright \text{Sub}(G)$.

Remark 5.6. Bass-Serre Theory asserts that if $\Lambda \curvearrowright T$ is a *minimal* action and if Λ is finitely generated, then the quotient graph $\Lambda \setminus T$ is compact. Indeed, by writing $\Lambda \setminus T$ as an increasing union of compact connected subgraphs κ_n , we can write G as the increasing union of the subgroups G_n corresponding to the graphs of groups of κ_n . Since G is finitely generated, this sequence of subgroups is stationary: $G_n = G_{n_0}$ for n greater than some n_0 . The pull-back of κ_{n_0} is a G -invariant cocompact subtree which coincides with T by minimality.

Question 5.7. What is the class of groups and minimal irreducible actions $G \curvearrowright T$ such that $\Lambda \leq G$ has infinite index if and only if $\Lambda \setminus T$ is infinite, i.e. for which

$$\text{Sub}_{|\bullet \setminus T|_\infty}(G) = \text{Sub}_{[\infty]}(G)?$$

Compare with Remarks 5.10 and 5.16, and with Proposition 5.25.

Thinking about this problem led us to the following question.

Question 5.8. Let H, A be infinite subgroups of the countable group G . Assume that A is amenable and that $\beta_1^{(2)}(G) > 0$ and $\beta_1^{(2)}(H) < \infty$. If $A \setminus G/H$ is finite, does this imply that H has finite index in G ?

Theorem 5.9 (A segment with finite stabiliser). *Assume $G \curvearrowright T$ is a minimal and irreducible action on a tree. Assume that there are two edges f_1, f_2 of T such that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite. Then the set $\text{Sub}_{|\bullet \setminus T|_\infty}(G)$ has no isolated point, thus its closure $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$ is a perfect set. In particular:*

$$\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} \subseteq \mathcal{K}(G).$$

Remark 5.10. Observe that $G \setminus T$ may be infinite in the above theorem and in this case

$$\text{Sub}_{|\bullet \setminus T|_\infty}(G) = \text{Sub}(G) = \mathcal{K}(G).$$

On the contrary, if $G \setminus T$ is finite, then $\text{Sub}_{|\bullet \setminus T|_\infty}(G) \subseteq \text{Sub}_{[\infty]}(G)$.

The subset $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$ is invariant under the conjugacy G -action. We will see that it moreover admits a natural finite partition into perfect G -invariant topologically transitive pieces. Indeed, recall from Proposition 3.2 that for any finite normal subgroup N there is

a finite clopen partition of $\text{Sub}(G)$ indexed by the set $\text{Conj}(N)^G$ of G -conjugacy classes of subgroups of N . This partition induces a G -invariant partition of $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$:

$$\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} = \bigsqcup_{C \in \text{Conj}(N)^G} \mathcal{F}_C^T$$

where, for each $C \in \text{Conj}(N)^G$, the subset

$$\mathcal{F}_C^T := \overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} \cap \{\Lambda \in \text{Sub}(G) : \Lambda \cap N \in C\} \quad (13)$$

is a perfect set by Theorem 5.9.

Theorem 5.11 (Topological transitivity). *Suppose that $G \curvearrowright T$ is a minimal and irreducible action on a tree. Assume that there are two edges f_1, f_2 of T such that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite, thus the kernel N_0 of the action $G \curvearrowright T$ is finite. Then*

- (i) *For each $C \in \text{Conj}(N_0)^G$ the action $G \curvearrowright \mathcal{F}_C^T$ is topologically transitive, and*
- (ii) *If $C \in \text{Conj}(N_0)^G$ has cardinality 1 then the action $G \curvearrowright \mathcal{F}_C^T$ is highly topologically transitive.*

In particular, if N_0 is trivial, then the action $G \curvearrowright \overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)}$ is highly topologically transitive.

Remark 5.12. The condition “ $C \in \text{Conj}(N_0)^G$ has cardinality 1” means that $M \in C$ is normalised by G ; it is satisfied for instance when N_0 is cyclic: a subgroup $M \leq N_0$ is uniquely defined by its cardinality. It is also satisfied for some finite index subgroup $G' \leq G$ that acts trivially on $\text{Sub}(N_0)$ by conjugation. Then G' satisfies Theorem 5.11(ii), but for a finer partition.

Remark 5.13 (Obstruction to topological 2-transitivity). Note that the condition is necessary in part (ii) of Theorem 5.11. Indeed, if C contains at least two distinct elements M, M' , then the action $G \curvearrowright \mathcal{F}_C^T$ is not topologically 2-transitive: Consider the four non-empty open sets $V_1 = V_2 = V_3 = \mathcal{F}_C^T \cap \{\Lambda \in \text{Sub}(G) : \Lambda \cap N_0 = M\}$ and $V_4 = \mathcal{F}_C^T \cap \{\Lambda \in \text{Sub}(G) : \Lambda \cap N_0 = M'\}$. For every element $\gamma \in G$ such that $\gamma V_1 \cap V_3 \neq \emptyset$, then $\gamma M \gamma^{-1} = M$ thus $\gamma V_2 \cap V_4 = \emptyset$.

Question 5.14. Do the conclusions of the above two theorems still hold if we replace the assumption about the edge stabilisers with the assumption that the kernel N_0 of $G \curvearrowright T$ is finite and the action of G/N_0 on the boundary of T is topologically free?

Proof of Theorem 5.9. Let us start by showing that, up to replacing f_2 with another edge, we can assume that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2) = N_0$, the kernel of $G \curvearrowright T$. In the connected component Z_2 of $T \setminus \{f_2\}$ that does not contain f_1 , let us choose an edge f_3 such that the intersection $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_3)$ has minimal cardinality; this is possible by the finiteness of $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ and because for every edge $f \in Z_2$, the edge f_2 belongs to the segment $[f_1, f]$. Let us call $P := \text{Stab}_G(f_1) \cap \text{Stab}_G(f_3)$. Let Z_3 be the connected component of $T \setminus \{f_3\}$ that does not contain f_1 . Every edge f in Z_3 satisfies

$$\text{Stab}_G(f_1) \cap \text{Stab}_G(f) = P. \quad (14)$$

By Lemma 5.2, there is an element $\gamma_0 \in G$ whose axis contains f_3 : a “half of the axis” of γ_0 is in Z_3 . Up to replacing γ_0 by its inverse, one can assume that $\gamma_0^{-1} \cdot f_3$ belongs to Z_3 .

For every edge e of T , there is a positive integer s such that $e \in \gamma_0^r \cdot Z_3$ for every $r \geq s$. Indeed, first note that $Z_3 \subseteq \gamma_0^r Z_3$ for all $r \geq 1$. Now if $e \notin Z_3$, let y be the projection of e on the axis of γ_0 , i.e. the vertex of $\text{Axis}(\gamma_0)$ that is the closest to $o(e)$. For every large enough positive power r of γ_0 , the vertex y lies between f_3 and $\gamma_0^r \cdot f_3$; and then y and e belong to $\gamma_0^{r+1} \cdot Z_3$. In particular, for every r larger than some s_0 , the segment $[f_1, f_3]$ is contained in $\gamma_0^r \cdot Z_3$. Every edge e in $\gamma_0^r \cdot Z_3$ satisfies $\text{Stab}_G(\gamma_0^r \cdot f_1) \cap \text{Stab}_G(e) = \gamma_0^r P \gamma_0^{-r}$, the γ_0^r -translated version of (14). In particular, for the edges f_1, f_3 and $r \geq s_0$: $\text{Stab}_G(\gamma_0^r \cdot f_1) \cap \text{Stab}_G(f_1) = \gamma_0^r P \gamma_0^{-r} = \text{Stab}_G(\gamma_0^r \cdot f_1) \cap \text{Stab}_G(f_3)$. It follows that $\gamma_0^r P \gamma_0^{-r} \leq \text{Stab}_G(f_1) \cap \text{Stab}_G(f_3) = P$. But P and $\gamma_0^r P \gamma_0^{-r}$ have the same finite cardinal, thus $P = \gamma_0^r P \gamma_0^{-r}$. It follows that P fixes every edge of T so that $P \leq N_0$. Since the kernel N_0 is contained in both $\text{Stab}_G(f_1)$ and $\text{Stab}_G(f_3)$, we also have $P \geq N_0$. We have proved that

$$\text{Stab}_G(f_1) \cap \text{Stab}_G(f_3) = N_0. \quad (15)$$

So, replace f_2 with f_3 .

If $\Lambda \in \text{Sub}_{|\bullet \setminus T|_\infty}(G)$ is not finitely generated, then Λ is a limit of finitely generated subgroups $\Lambda_n \leq \Lambda$ which satisfy a fortiori that $\Lambda_n \setminus T$ is infinite. This is the case, in particular, if $G \setminus T$ is not finite: by minimality, G is not finitely generated.

It is thus enough to prove that $\Lambda \in \mathcal{K}(G)$ when $\Lambda \in \text{Sub}_{|\bullet \setminus T|_\infty}(G)$ is finitely generated, by showing that Λ satisfies the conditions of Theorem 5.5. Since Λ is finitely generated, it admits a cocompact (thus proper) invariant subtree $T_\Lambda \subsetneq T$ (see Remark 5.6).

Claim 1. *Let f be any edge of $T \setminus T_\Lambda$ and let Z_f be the connected component of $T \setminus \{f\}$ that does not contain T_Λ . Let L be the segment $[f_1, f_2]$. One can assume, up to replacing L with some G -translate, that $L \subseteq Z_f$ and that f_1 separates f_2 from f and T_Λ .*

Proof of Claim 1. Since N_0 is normal, for any $\gamma \in G$ the pair $(\gamma \cdot f_1, \gamma \cdot f_2)$ satisfies the same assumption as (f_1, f_2) . Then Claim 1 is a direct application of the fourth item of Lemma 5.2 (existence of γ_2). \diamond

Claim 2. *Suppose that f, Z_f and L are as in the Claim 1 and define the edge $e_\Lambda := f_2$. Then*

$$\text{Stab}_\Lambda(e_\Lambda) = \Lambda \cap N_0. \quad (16)$$

Proof of Claim 2. If λ belongs to $\text{Stab}_\Lambda(e_\Lambda)$, then (since T_Λ is Λ -invariant) the whole path from e_Λ to T_Λ must be point-wise fixed by λ . This path uses the subpath L whose edges f_1 and f_2 satisfy $\text{Stab}_\Lambda(f_1) \cap \text{Stab}_\Lambda(f_2) \leq \text{Stab}_G(f_1) \cap \text{Stab}_G(f_2) = N_0$. Thus $\lambda \in \Lambda \cap N_0$, i.e. $\text{Stab}_\Lambda(e_\Lambda) \leq \Lambda \cap N_0 \leq \text{Stab}_\Lambda(e_\Lambda)$ since N_0 is the kernel of the action. Claim 2 is proved. \diamond

We have proved that Λ (when finitely generated) satisfies the conditions of Theorem 5.5, i.e. $\Lambda \in \mathcal{N}_T$, in the terminology of that theorem. It follows that $\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} \subseteq \overline{\mathcal{N}_T}$. Observe that for every $\Lambda' \in \mathcal{N}_T$, the quotient $\Lambda' \setminus T$ is infinite (one of the components of $T \setminus \{e_{\Lambda'}\}$ is sent injectively in $\Lambda' \setminus T$), thus $\mathcal{N}_T \subseteq \text{Sub}_{|\bullet \setminus T|_\infty}(G)$. Hence

$$\overline{\text{Sub}_{|\bullet \setminus T|_\infty}(G)} = \overline{\mathcal{N}_T}. \quad (17)$$

Recall from Theorem 5.5 that \mathcal{N}_T has no isolated points with respect to its induced topology. This concludes the proof of Theorem 5.9. \square

We now prove Theorem 5.11, keeping the same notation and numbering of claims.

Proof of Theorem 5.11. Fix $C \in \text{Conj}(N_0)^G$.

In order to prove (i), we will have to show that if V_1, V_2 are two arbitrary non-empty open sets of \mathcal{F}_C^T , then there is $\gamma \in G$ such that $\gamma V_1 \gamma^{-1} \cap V_2 \neq \emptyset$.

Finitely generated subgroups are dense in $\text{Sub}_{|\bullet \setminus T|^\infty}(G)$ and $\{\Lambda \in \text{Sub}(G) : \Lambda \cap N_0 \in C\}$ is open, so there exist finitely generated $\Lambda_i \in V_i$ for $i = 1, 2$. By Theorem 5.9, $\text{Sub}_{|\bullet \setminus T|^\infty}(G)$ has no isolated points, so we can assume that $\Lambda_1, \Lambda_2 \notin \text{Sub}(N_0)$. Moreover, we can assume that $\Lambda_1 \cap N_0 = \Lambda_2 \cap N_0 =: M$. Indeed, if $M_i := \Lambda_i \cap N_0$ for $i = 1, 2$ then there exists $\gamma_2 \in G$ such that $\gamma_2 M_2 \gamma_2^{-1} = M_1$, and we can replace V_2 by $V_2' := \gamma_2 V_2 \gamma_2^{-1}$: if there exists γ such that $\gamma V_1 \gamma^{-1} \cap V_2' \neq \emptyset$ then $\gamma_2^{-1} \gamma V_1 \gamma^{-1} \gamma_2 \cap V_2 \neq \emptyset$.

To prove part (ii), we need to show that, if $C = \{M\}$ (i.e. M is normal in G) then, for all positive integers r and for all non-empty open sets $V_1, \dots, V_{2r} \subseteq \mathcal{F}_C$, there exists $\gamma \in G$ such that $\gamma V_i \cap V_{r+i} \neq \emptyset$ for each $i \in \{1, \dots, r\}$. As above, we can pick $\Lambda_i \in V_i$ that is finitely generated and not contained in $\text{Sub}(N_0)$. Moreover $\Lambda_i \cap N_0 = M$ for each i by definition. Thus both part (i) and part (ii) follow from the following statement:

Fix $C \in \text{Conj}(N_0)^C$ and $M \in C$ (we don't assume that $|C| = 1$). Let $r \geq 1$ and $V_1, \dots, V_{2r} \subseteq \mathcal{F}_C$ be open subsets such that, for each $i \in \{1, \dots, 2r\}$, there exist finitely generated subgroups $\Lambda_i \in V_i$ where $\Lambda_i \cap N_0 = M$ and $\Lambda_i \neq M$. Then there is an element $\gamma \in G$ such that $\gamma V_i \cap V_{r+i} \neq \emptyset$ for each $i \in \{1, \dots, r\}$.

Let us prove that this statement indeed holds. For any subgroup $H \leq G$ let T_H denote a minimal H -invariant subtree of T . Recall that either T_H is a single vertex or T_H is the union of the axes of the hyperbolic elements of H .

Claim 3. *If T_H contains a half-tree, i.e. a connected component Z of $T \setminus \{e\}$ for some edge $e \in E(T)$, then $T_H = T$. If H is moreover finitely generated, then $H \setminus T$ is finite.*

Proof of Claim 3. Let f be an edge in Z and $h \in H$ a hyperbolic element whose axis $\text{Axis}(h)$ contains f . Then, up to replacing h by h^{-1} , the image $h \cdot e$ belongs to Z . Let $g' \in H$ be a hyperbolic element whose axis contains $h \cdot e$. Then the axis of the conjugate $g = h^{-1} g' h \in H$ contains e . Thus $e \in T_H$ and T_H contains all the half-trees $g^k(Z)$ for $k \in \mathbb{Z}$. It follows that $T = \cup_k g^k(Z) = T_H$. When H is finitely generated, then (Remark 5.6) there is a finite connected subgraph $W \subseteq H \setminus T$ whose associated graph of groups has fundamental group equal to H . If $H \setminus T$ is infinite, then the pull-back of W in T is a proper H -invariant subtree of T . This proves Claim 3. \diamond

Claim 4. *Assume that $G \curvearrowright T$ is a minimal and irreducible action on a tree and that $T_i := T_{\Lambda_i} \subsetneq T$ are proper Λ_i -invariant subtrees for finitely generated subgroups $\Lambda_i \leq G$ for $i \in \{1, \dots, 2r\}$. Then, there is an edge $e \in E(T)$ such that $\cup_{i=1}^{2r} T_i$ is contained in one connected component of $T \setminus \{e\}$ and such that $\text{Stab}_{\Lambda_i}(e) = M$ for all i .*

Proof of Claim 4. If U, V are subtrees of T that do not contain any half-tree, then the convex hull of $U \cup V$ does not contain any half-tree. To see this, pick an edge $e_0 \in E(T)$ and let K be one of its half-trees. Since U does not contain K nor any half-tree therein, there is an edge $e_1 \in E(K)$ such that U is in the same half-tree of $T \setminus \{e_1\}$ as e_0 . Let K_2 be the other half-tree of $T \setminus \{e_1\}$ and note that $K_2 \subseteq K$. Since no half-tree is contained in V , there exists an edge $e_2 \in E(K_2)$ such that V is in the same half-tree of $T \setminus \{e_2\}$ as e_1 (and e_0). If $K_3 \subseteq T \setminus \{e_2\}$

is the half-tree which does not contain e_1 then $K_3 \subseteq K$ and K_3 has empty intersection with the convex hull of $U \cup V$. Thus the half-tree K is not entirely contained in the convex hull of $U \cup V$.

Since minimal subtrees T_i of Λ_i do not contain any half-tree by Claim 3, the same holds for their convex hull. It follows by induction that $\cup_{i=1}^{2r} T_i$ is contained in a half-tree of $T \setminus \{f\}$ for some $f \in E(T)$. This remains true if we replace f by any edge in the connected component of $T \setminus \{f\}$ which does not contain $\cup_{i=1}^{2r} T_i$.

By Claim 1, one can assume that the segment $L = [f_1, f_2]$ is contained in the connected component of $T \setminus \{f\}$ that does not contain $\cup_{i=1}^{2r} T_i$ and that f_1 separates f_2 from f , thus from each T_i . By Claim 2, the edge $e = f_2$ satisfies $\text{Stab}_{\Lambda_i}(e) = \Lambda_i \cap N_0 = M$ for each i and this concludes the proof of Claim 4. \diamond

We take an edge $e \in E(T)$ as in the Claim 4 and let Z be the connected component of $T \setminus \{e\}$ that does not contain $\cup_{i=1}^{2r} T_i$. By Lemma 5.2, there is a hyperbolic element $g \in G$ whose axis is contained in Z .

We now fix $i \in \{1, \dots, r\}$ and show that Λ_i satisfies the assumptions of Lemma 5.4 where the Λ_i -invariant proper subtree is $T_{\Lambda_i} = T_i$ and the subgroups H_n are of the form $g^q \Lambda_{r+i} g^{-q}$. The sequences of subtrees $(S_n)_n$ and of edges $(e_n)_n$ will be defined shortly.

Since the action by conjugation of G on $\text{Sub}(N_0)$ has finite orbits, there is a non-zero integer k such that $g^k M g^{-k} = M$. It follows that $g^{kn} \Lambda_{r+i} g^{-kn} \cap N_0 = g^{kn} (\Lambda_{r+i} \cap N_0) g^{-kn} = g^{kn} M g^{-kn} = M$ for every integer n .

Let $e_n := g^{kn} \cdot e$. Then the distance of e_n from T_{Λ_i} is greater than $n + 1$ times the translation length of g . The subtree $S_n := g^{kn} \cdot T_{r+i}$ is contained in Z and e_n separates S_n from T_i . The subtree $g^{kn} \cdot T_{r+i}$ is an invariant subtree for the subgroup $H_n := g^{kn} \Lambda_{r+i} g^{-kn}$. We have defined (H_n) , (S_n) and (e_n) and checked Assumptions 1, 2, 3a and 3b of Lemma 5.4. The assumption we made that Λ_{r+i} is not a subgroup of N_0 implies that $\text{Stab}_{H_n}(e_n) \neq H_n$, thus checking Assumption 3d. If an element $\lambda \in \Lambda_i$ sends some e_n to an edge $\lambda \cdot e_n$ belonging to the path from some other e_k to T_i then λ must fix the projection p of e_k in T_i and $\lambda \cdot e_n$ belongs to Z . It follows that λ must fix the edge e and thus $\lambda \in \text{Stab}_{\Lambda_i}(e) = M \leq N_0$. So $\Lambda_i \cdot e_n$ does not separate e_k from the tree T_i . Therefore, (e_n) verifies condition 3e of Lemma 5.4. Concerning the stabilisers: $\text{Stab}_{\Lambda_i}(e_n) = g^{kn} \text{Stab}_{\Lambda_i}(e) g^{-kn} = g^{kn} M g^{-kn} = M$ while $\text{Stab}_{H_n}(e_n) = \text{Stab}_{g^{kn} \Lambda_{r+i} g^{-kn}}(g^{kn} \cdot e) = \text{Stab}_{\Lambda_{r+i}}(e) = M$. The Assumption 3c is satisfied.

Thus, by Lemma 5.4 (I) and (II), $\Delta_n^i := \langle \Lambda_i, g^{kn} \Lambda_{r+i} g^{-kn} \rangle = \Lambda_i *_M g^{kn} \Lambda_{r+i} g^{-kn}$ and we have the non-trivial convergence:

$$\Delta_n^i = \Lambda_i *_M g^{kn} \Lambda_{r+i} g^{-kn} \xrightarrow[n \rightarrow \infty]{} \Lambda_i.$$

Since $\{\Lambda \leq G : \Lambda \cap N_0 = M\}$ is open, we have $\Delta_n^i \cap N_0 = \Lambda_i \cap N = M$ for all large enough n . By (V) of Lemma 5.4, the quotient $\Delta_n^i \backslash T$ is infinite, i.e. $\Delta_n^i \in \text{Sub}_{|\bullet| \backslash T|_\infty}(G)$. Thus $\Delta_n^i \in \mathcal{F}_C^T$. It follows that for n larger than some n_i , the subgroup $\Delta_n^i \in V_i$.

It remains to show that we have the non-trivial convergence:

$$g^{-kn} \Delta_n^i g^{kn} = g^{-kn} \Lambda_i g^{kn} *_M \Lambda_{r+i} \xrightarrow[n \rightarrow \infty]{} \Lambda_{r+i}.$$

It is enough to observe that $g^{-kn} \Delta_n^i g^{kn} = g^{-kn} \Lambda_i g^{kn} *_M \Lambda_{r+i}$ is obtained by the exact symmetric construction by exchanging the roles of Λ_i, Λ_{r+i} and replacing g by g^{-1} . So the

symmetric application of Lemma 5.4 (II) shows that $\lim_{n \rightarrow \infty} g^{-kn} \Delta_n^i g^{kn} = \Lambda_{r+i}$. It follows as above that for all n larger than some n_{r+i} , the subgroup $g^{-kn} \Delta_n^i g^{kn} \in V_{r+i}$.

Hence, for any $n \geq \max\{n_j : j \in \{1, \dots, 2r\}\}$ and for each $i \in \{1, \dots, r\}$, we have:

$$\Delta_n^i \in V_i \text{ and } g^{-kn} \Delta_n^i g^{kn} \in V_{r+i}, \quad \text{and so } g^{-kn} V_i g^{kn} \cap V_{r+i} \neq \emptyset.$$

This completes the proof of Theorem 5.11. □

5.3 Applications

Recall that the G -action $G \curvearrowright T$ on a tree is **acylindrical** if there is $R > 0$ such that the stabilisers of any path of length at least R is trivial.

Examples of 3-acylindrical actions include the actions on the Bass-Serre tree of non-trivial amalgamated free products $A *_C B$ where C is **malnormal** in A (i.e. for all $g \in A \setminus C$, $C \cap gCg^{-1}$ is trivial).

Corollary 5.15 (Acylindrical actions). *Let $G \curvearrowright T$ be a minimal irreducible and acylindrical action of a countable group on a tree. If $\Lambda \leq G$ is a subgroup such that the quotient graph $\Lambda \backslash T$ is infinite, then $\Lambda \in \mathcal{K}(G)$. Moreover, $G \curvearrowright \overline{\text{Sub}_{|\bullet|_T|_\infty}(G)}$ is highly topologically transitive.*

Proof. The G -action on T satisfies the assumptions of Theorem 5.9 and Theorem 5.11: the intersection of the stabilisers of any pair of edges at distance greater than the acylindricity constant R is trivial. □

Remark 5.16. There exist acylindrical actions $G \curvearrowright T$ as in Corollary 5.15 where G admits a finitely generated subgroup $\Lambda \leq G$ with infinite index but where the quotient $\Lambda \backslash T$ is finite. Moreover, there are examples where Λ belongs to $\mathcal{K}(G)$ and others where it does not.

For example, let C be any finitely generated infinite torsion-free group and let A, B be finitely generated non-trivial groups such that C is malnormal in $G_1 = A \rtimes C$ and $G_2 = B \rtimes C$ respectively. Such groups exist by D. Osin's Appendix in [HW14, Theorem 15].

Then let $G = G_1 *_C G_2$ where the injections $C \hookrightarrow G_i$ are given by the natural inclusion maps and consider the action of G on the Bass-Serre tree T associated to this decomposition. It follows from the malnormality of C in G_1 and G_2 that the action $G \curvearrowright T$ is 2-acylindrical. Moreover, the normal subgroup $\Lambda = \langle A, B \rangle \trianglelefteq G$ is finitely generated, has infinite index in G since $G/\langle A, B \rangle \simeq C$, and acts transitively on the set of edges of T , so $\Lambda \backslash T$ consists of a single edge.

Moreover, by Proposition 3.5, the finitely generated normal subgroup $\Lambda = \langle A, B \rangle = A * B$ belongs to the perfect kernel $\mathcal{K}(G)$ if and only if the trivial group $\{\text{id}\}$ belongs to the perfect kernel $\mathcal{K}(C)$. If we choose for instance $C = \mathbb{Z}$ then $\Lambda \notin \mathcal{K}(G)$ while if $C = \mathbf{F}_2$, then $\Lambda \in \mathcal{K}(G)$.

The results of the previous section imply that:

If G is a finitely generated non-elementary free product¹ then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ and the action of G on its perfect kernel is highly topologically transitive. More generally,

Corollary 5.17 (Infinitely many ends). *If G is a finitely generated group with infinitely many ends, then the perfect kernel of G consists of its subgroups of infinite index:*

$$\mathcal{K}(G) = \text{Sub}_{[\infty]}(G).$$

¹A free product $G = A * B$ is **non-elementary** if $|A|, |B| \geq 2$ and at least one of A and B has cardinality at least 3.

Moreover, G has a greatest finite normal subgroup N_0 and the set $\text{Conj}(N_0)^G$ defines a finite clopen partition of $\mathcal{K}(G)$ into G -invariant and topologically transitive perfect subsets:

$$\mathcal{K}(G) = \bigsqcup_{C \in \text{Conj}(N_0)^G} \mathcal{S}_C, \quad \text{where } \mathcal{S}_C := \text{Sub}_{[\infty]}(G) \cap \{\Lambda : \Lambda \cap N_0 \in C\}.$$

If $N_0 = \{\text{id}\}$, resp. $|C| = 1$, then $G \curvearrowright \mathcal{K}(G)$, resp. $G \curvearrowright \mathcal{S}_C$, is highly topologically transitive.

Proof. By Stallings' theorem [Sta68, Sta71], G admits a minimal and irreducible action $G \curvearrowright T$ on a tree T with finite edge stabilisers. If M is a finite normal subgroup of G , then it is contained in the kernel of the action $G \curvearrowright T$. Conversely, this kernel is finite and normal: this is N_0 . If $\Lambda \in \text{Sub}(G)$ has infinite index, then the image in $\Lambda \backslash T$ of the G -orbit any edge e of T is in bijection with the double classes $\Lambda \backslash G / \text{Stab}_G(e)$, which is infinite since $\text{Stab}_G(e)$ is finite. Thus $\Lambda \backslash T$ is infinite and \mathcal{S}_C coincides with \mathcal{F}_C^T (defined in Equation 13). Since the edge stabilisers are finite, any two distinct edges f_1, f_2 of T satisfy a fortiori that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite. We conclude by applying Theorem 5.9 and Theorem 5.11. \square

Example 5.18 ($\text{SL}(2, \mathbb{Z})$). This corollary applies for instance to $G = \text{SL}(2, \mathbb{Z})$ for which $N_0 = \{\pm \text{id}\} \simeq \mathbb{Z}/2\mathbb{Z}$ is the centre. The perfect kernel decomposes into two invariant pieces

$$\mathcal{K}(G) = \text{Sub}_{[\infty]}(\text{SL}(2, \mathbb{Z})) = \{\Lambda : N_0 \leq \Lambda\} \sqcup \{\Lambda : N_0 \not\leq \Lambda\}$$

on which the action is highly topologically transitive.

Corollary 5.19. *The action of the free group $\mathbf{F}_{\mathbb{N}}$ on countably many generators on its perfect kernel $\mathcal{K}(\mathbf{F}_{\mathbb{N}}) = \text{Sub}(\mathbf{F}_{\mathbb{N}})$ is highly topologically transitive.*

Proof. By Proposition 3.8, $\mathcal{K}(\mathbf{F}_{\mathbb{N}}) = \text{Sub}(\mathbf{F}_{\mathbb{N}})$. Every non-empty open subset V of $\text{Sub}(\mathbf{F}_{\mathbb{N}})$ contains a finitely generated subgroup, thus for every $2r$ -tuple $(V_1, V_2, \dots, V_{2r})$ of non-empty open sets, there is a finite subset $J \subseteq \mathbb{N}$ with $|J| \geq 2$ such that $V_i \cap \text{Sub}_{[\infty]}(\mathbf{F}_J) \neq \emptyset$ for all i . By Corollary 5.17, there is $g \in \mathbf{F}_J$ such that $g \cdot (V_i \cap \text{Sub}_{[\infty]}(\mathbf{F}_J)) \cap (V_{r+i} \cap \text{Sub}_{[\infty]}(\mathbf{F}_J)) \neq \emptyset$ for all i ; in particular, $g \cdot V_i \cap V_{r+i} \neq \emptyset$. \square

Example 5.20 (Weakly malnormal edge group). A subgroup $C \leq G$ is said to be **weakly malnormal** if there exists $g \in G$ such that $gCg^{-1} \cap C$ is finite. Let G be a free product with amalgamation $G = A *_C B$ or a HNN-extension $G = A *_C$ with irreducible associated Bass-Serre tree T . Assume that C is weakly malnormal in G . It follows that the action of G on T verifies the conditions of Theorem 5.9 (and of Theorem 5.11). Thus every subgroup Λ such that $\Lambda \backslash T$ is infinite belongs to $\mathcal{K}(G)$.

For instance, if $A = \mathbf{F}_p$ is a non-abelian free group, then any finitely generated subgroup C with infinite-index is weakly malnormal in A and thus in $G = A *_C B$ for any B . More generally, by [PT11, Theorem 5.12] (with $\beta_1^{(2)}(\cdot)$ the first ℓ^2 -Betti numbers): *If $C \leq G$ is an infinite index subgroup with $\beta_1^{(2)}(C) < \infty$ and if $\beta_1^{(2)}(G) > 0$, then C is weakly malnormal.*

This leads to plenty of examples. Take $A \geq C$ and $B \geq C$ such that $[A : C] \geq 2$ and $[B : C] \geq 3$: this will ensure the irreducibility of $G \curvearrowright T$ by Proposition 5.1. Assume that $\beta_1^{(2)}(C) < \infty$ and $\beta_1^{(2)}(A) + \beta_1^{(2)}(B) - \beta_1^{(2)}(C) > 0$: an application of the Mayer-Vietoris formula from Cheeger-Gromov [CG86, p. 204, §4] gives (when C is infinite) $\beta_1^{(2)}(A *_C B) \geq \beta_1^{(2)}(A) + \beta_1^{(2)}(B) - \beta_1^{(2)}(C)$. Thus C is weakly malnormal in $G = A *_C B$ and Theorems 5.9 and 5.11 apply.

5.4 A rich class of groups with maximal perfect kernel

In their study of the subgroups of direct products of limit groups, Bridson, Howie, Miller, and Short [BHMS09] introduced a class \mathcal{C} of finitely presented groups defined in a hierarchical manner as the union of classes \mathcal{C}_n and proved a series of properties of this class, in particular their Proposition 5.2. A close look at their proof reveals that various properties that are of interest to us are verified by the larger class \mathcal{Q} defined below, with essentially the same arguments. Compare with [BHMS09, Section 2.2]: \mathcal{C}_0 is the class of non-elementary free products of finitely presented groups, and a group lies in \mathcal{C}_n if and only if it is the fundamental group of a finite, acylindrical graph of finitely presented groups, where all of the edge groups are cyclic, and at least one of the vertex groups lies in \mathcal{C}_{n-1} .

Definition 5.21 (The classes \mathcal{Q} and \mathcal{Q}'). The class \mathcal{Q}_0 consists of finitely generated groups with infinitely many ends. A group lies in \mathcal{Q}_n if it is the fundamental group of a finite graph of groups such that

- every vertex group is finitely generated,
- every edge group is finitely generated virtually abelian,
- at least one vertex group lies in \mathcal{Q}_{n-1} and
- the associated Bass-Serre tree T admits a pair (f_1, f_2) of edges whose stabilisers have finite intersection.

The class \mathcal{Q} is the union of the classes \mathcal{Q}_n . The class \mathcal{Q}' is the subclass of groups in \mathcal{Q} without any non-trivial finite normal subgroup.

By Stallings' Theorem a group in \mathcal{Q}_0 admits a minimal irreducible action on a tree with a single orbit of edges and with finite edge stabilisers. We call any such G -tree a “Stallings tree” of G . A G -tree T witnessing that $G \in \mathcal{Q}$ will be called “the” Bass-Serre tree of G , although it is not uniquely defined.

Observe that the class \mathcal{Q} does not contain any infinite virtually abelian group as all its elements contain a non-abelian free subgroup.

Example 5.22. The class \mathcal{Q}' contains all non-abelian limit groups ([BHMS09, Corollary 2.2]). Another important class of groups contained in \mathcal{Q}' is that of graphs of free groups with cyclic edges such that at least one vertex group is non-cyclic.

The following is a corollary of Theorems 5.9 and 5.11.

Corollary 5.23. *Let G be any group in the class \mathcal{Q} . Then $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$. Moreover, the conclusions of Theorem 5.11 apply. In particular, if $G \in \mathcal{Q}'$, then the action $G \curvearrowright \mathcal{K}(G)$ is highly topologically transitive.*

We postpone the proof of this corollary until after the following two necessary propositions and after their proofs (which mimic that of [BHMS09, Proposition 5.2]). Observe that the assumptions of the propositions both repeat those defining the class \mathcal{Q} each with one condition missing (the third and the fourth respectively).

Proposition 5.24. *Let G be the fundamental group of a finite graph of finitely generated groups, where all of the edge groups are finitely generated virtually abelian and the associated Bass-Serre tree T admits a pair (f_1, f_2) of edges whose stabilisers have finite intersection. Let $\Lambda, C \leq G$ be finitely generated subgroups where C is virtually abelian. If the set of double classes $\Lambda \backslash G / C$ is finite, then the quotient graph $\Lambda \backslash T$ is finite.*

This proposition applies in particular to any $G \in \mathcal{Q}$ and its Bass-Serre tree.

Proposition 5.25. *Let G be the fundamental group of a finite graph of finitely generated groups, where all of the edge groups are virtually abelian. Assume that the associated Bass-Serre tree T admits a vertex w satisfying $\text{Stab}_G(w) \in \mathcal{Q}_n$, for $n \geq 0$. Then the quotient graph $\Lambda \backslash T$ is finite if and only if the index $[G : \Lambda]$ is finite.*

Proof of Proposition 5.24. We consider two cases.

Case 1. Suppose that C fixes a vertex v of T . Then, by the double-coset hypothesis, the G -orbit of v consists of only finitely many Λ -orbits $\Lambda \cdot v_i$. Since the action of G on T is cocompact, every point in T is at a bounded distance from $G \cdot v$, thus $\Lambda \backslash T$ has finite diameter. The fundamental group of the topological graph underlying $\Lambda \backslash T$ is finitely generated since Λ is finitely generated and maps onto it. By minimality of $G \curvearrowright T$, there are no valency 1 edges in T . Moreover, the valency of a vertex z in the tree T is the sum of the indices $[\text{Stab}_\Lambda(z) : \text{Stab}_\Lambda(e)]$ over a subset of the edges e incident to z containing a representative from each Λ -orbit. It follows that if z is a vertex with $\text{Stab}_\Lambda(z) = \text{Stab}_\Lambda(e)$ for some incident edge e then the image of z in $\Lambda \backslash T$ does not have valency 1. By finite generation of Λ , only finitely many valency 1 vertices of $\Lambda \backslash T$ can have a vertex group larger than the group of their incident edge. It follows that $\Lambda \backslash T$ is finite.

Case 2. Suppose that C does not fix a vertex. Since C is finitely generated, it admits a hyperbolic element $c \in C$ [Ser80, Corollary 3, p. 65]. Let A be the axis of c . Let $C' \leq C$ be a finite index abelian subgroup. For some integer j , we have $c' := c^j \in C'$, and A is also the axis of c' . Let $a \in C$. The image $a \cdot A$ is the axis of the hyperbolic element $ac'a^{-1}$. In case $a \in C'$ then $ac'a^{-1} = c'$, so the line A is C' -invariant and A is the axis of all hyperbolic elements of C' . If $a \in C$, a certain power k satisfies $(aca^{-1})^k \in C'$, thus the axis of aca^{-1} is also A . It follows that A is C -invariant and is the minimal subtree of C .

Let $e \in E(T)$ be an edge that belongs to A . Contract to one point each connected component of $T \setminus G \cdot e$ in order to produce the G -equivariant factor map $P : T \rightarrow S$ to the edge-transitive G -tree S . Up to enlarging the segment $L := [f_1, f_2]$ of T so as to start and finish with edges in $G \cdot e$, the image $P(L) = [P(f_1), P(f_2)]$ satisfies $|\text{Stab}_G(P(L))| < \infty$. Let $\bar{e} := P(e)$. The minimal tree of C in S is the image $\bar{A} := P(A) = C \cdot \bar{e}$ of the axis A . The two maps $(G \rightarrow E(S); g \mapsto g \cdot \bar{e})$ and $(\Lambda \backslash G \rightarrow E(\Lambda \backslash S); \Lambda g \mapsto \Lambda(g \cdot \bar{e}))$ are onto and the finiteness of the double cosets $\Lambda \backslash G / C$ thus implies the following:

Claim: The quotient graph $\Lambda \backslash S$ is covered by finitely many images $\Lambda g_i C \cdot \bar{e}$, i.e. by finitely many images $Q_\Lambda(g_i(\bar{A}))$ of lines under the quotient map $Q_\Lambda : S \rightarrow \Lambda \backslash S$. \diamond

Let $S_\Lambda \subseteq S$ be a minimal Λ -invariant subtree. Assume for a contradiction that $S_\Lambda \neq S$. Let ε be an edge of $S \setminus S_\Lambda$ and Z_1 the half-tree of $S \setminus \{\varepsilon\}$ that does not contain S_Λ . By Lemma 5.2, there is $\gamma_2 \in G$ such that $\gamma_2([P(f_1), P(f_2)]) \subseteq Z_1$ and $\gamma_2(P(f_1))$ separates $\gamma_2(P(f_2))$ from ε . We show that the map Q_Λ is boundedly finite-to-one in restriction to the half-tree Z_2 of $S \setminus \{\gamma_2(P(f_2))\}$ that does not contain S_Λ . Indeed an element $\lambda \in \Lambda$ sending a vertex u of Z_2 inside Z_2 must fix the projection v of u on S_Λ and thus the segment

$[v, \gamma_2(P(f_2))]$. But this segment contains the segment $\gamma_2(P(L))$. Thus λ belongs to the γ_2 -conjugate $\gamma_2 \text{Stab}_G(P(L))\gamma_2^{-1}$ of the point-wise stabiliser of $P(L)$, which is finite. The map $Z_2 \rightarrow Q_\Lambda(Z_2) = (\gamma_2 \text{Stab}_G(P(L))\gamma_2^{-1} \cap \Lambda) \backslash Z_2$ has fibers of uniformly bounded size, thus $Q_\Lambda(Z_2)$ has infinitely many ends. The graph $\Lambda \backslash S$ contains $Q_\Lambda(Z_2)$, thus it has infinitely many ends, and it cannot be covered by finitely many images of lines, contrarily to the above Claim. Thus $S_\Lambda = S$.

Since Λ is finitely generated, $\Lambda \backslash S$ is finite (Remark 5.6). It follows that $\Lambda \backslash T$ is finite. Proposition 5.24 is proved. \square

Proof of Proposition 5.25. We will argue by induction on n .

The initialisation of the induction is that groups in \mathcal{Q}_0 satisfy the conclusion of Proposition 5.25: If $G \in \mathcal{Q}_0$, i.e. its Stallings tree is an irreducible edge transitive action $G \curvearrowright T$ with a finite edge stabiliser K , then for any $\Lambda \leq G$, the quotient $\Lambda \backslash T$ is finite if and only if $\Lambda \backslash G/K$ is finite, if and only if $\Lambda \backslash G$ is finite.

Assume the conclusion of Proposition 5.25 holds for all groups satisfying the assumptions up to some $n - 1$. In particular, this conclusion holds for all groups in \mathcal{Q}_n .

Let now G be a group satisfying the assumptions for n with $\text{Stab}_\Lambda(w) \in \mathcal{Q}_n$. Let $\Lambda \leq G$ and assume $\Lambda \backslash T$ is finite. Observe that $\text{Stab}_\Lambda(w)$ is also finitely generated since (a) subgroups of finitely generated (virtually) abelian groups are finitely generated and (b) a finitely generated graph of groups (such as Λ) with finitely generated edge groups must have finitely generated vertex groups by [Coh89, p.43, Propositions 29 and 35].

Let e be some edge incident at w . Then $|\text{Stab}_\Lambda(w) \backslash \text{Stab}_G(w) / \text{Stab}_G(e)|$ is bounded above by the finite number of edges of $\Lambda \backslash T$ that are incident at $\Lambda \cdot w \in \Lambda \backslash T$. Since $\text{Stab}_G(w) \in \mathcal{Q}_n$, since $\text{Stab}_\Lambda(w)$ and $\text{Stab}_G(e)$ are finitely generated and $\text{Stab}_G(e)$ is virtually abelian, then applying Proposition 5.24 to the action of $\text{Stab}_G(w)$ on its own Bass-Serre tree T' implies that $\text{Stab}_\Lambda(w) \backslash T'$ is finite. Then, by the induction hypothesis (including the initialisation when $n = 0$), $\text{Stab}_\Lambda(w)$ has finite index in $\text{Stab}_G(w)$. And similarly, for each $g \in G$, the subgroup $\text{Stab}_\Lambda(gw)$ has finite index in $\text{Stab}_G(gw)$: Indeed $\text{Stab}_G(gw) \simeq \text{Stab}_G(w) \in \mathcal{Q}_n$ and gw could have played the role of w .

Consider the action of $\text{Stab}_G(w)$ by right multiplication on $\Lambda \backslash G$: the orbits are the double cosets $\Lambda \backslash G / \text{Stab}_G(w)$ and hence are finite in number because they index a subset of the vertices of $\Lambda \backslash T$; moreover, the stabilizer of Λg is $g^{-1} \Lambda g \cap \text{Stab}_G(w) = g^{-1} (\Lambda \cap g \text{Stab}_G(w) g^{-1}) g = g^{-1} \text{Stab}_\Lambda(gw) g$, which we have just seen has finite index in $g^{-1} \text{Stab}_G(gw) g = \text{Stab}_G(w)$. Thus $\Lambda \backslash G$ is finite. The converse is clear: if $\Lambda \backslash G$ is finite, the quotient $\Lambda \backslash T$ is finite since $G \backslash T$ is finite. Proposition 5.25 is proved. \square

Proof of Corollary 5.23. An element $G \in \mathcal{Q}$ is equipped with its defining Bass-Serre G -tree T . Observe that the action $G \curvearrowright T$ can be assumed to be minimal and irreducible: take the minimal j such that $G \in \mathcal{Q}_j$ and note that the stabiliser of w is not virtually abelian, thus the stabiliser of any edge incident to w has infinite index in $\text{Stab}_G(w)$. Irreducibility of the action then follows Proposition 5.1. By Theorem 5.9 $\text{Sub}_{|\bullet \backslash T|_\infty}(G) \subseteq \mathcal{K}(G) \subseteq \text{Sub}_{[\infty]}(G)$. It is enough to check that $\overline{\text{Sub}_{|\bullet \backslash T|_\infty}(G)} \supseteq \text{Sub}_{[\infty]}(G)$.

If $\Lambda \in \text{Sub}_{[\infty]}(G)$ is finitely generated then $\Lambda \backslash T$ is infinite: if $n = 0$ this is because the action of G on the edges of its Stallings tree is transitive with finite stabilisers, and otherwise this follows from Proposition 5.25. Thus $\Lambda \in \text{Sub}_{|\bullet \backslash T|_\infty}(G)$. If $\Lambda \in \text{Sub}_{[\infty]}(G)$ is not finitely generated, it is a limit of proper finitely generated subgroups $\Lambda_n \leq \Lambda$, which are thus also of

infinite index, so that $\Lambda_n \in \text{Sub}_{|\bullet| \setminus T|_\infty}(G)$ and $\Lambda \in \overline{\text{Sub}_{|\bullet| \setminus T|_\infty}(G)}$. Then Theorem 5.11 applies concerning the dynamics. \square

6 Hyperbolic groups

6.1 Generalities on the space of subgroups of hyperbolic groups

Recall from [Gro87, 5.3] that a subset Y of a geodesic metric space X is **quasiconvex** if its convex hull is contained within a bounded neighbourhood of Y .

Let G be a hyperbolic (aka word-hyperbolic) group. A subgroup $H \leq G$ is **quasiconvex** if it is a quasiconvex subset of the Cayley graph X of G associated with a finite generating set of G . Note that this does not depend on the choice of Cayley graph as quasiconvexity in hyperbolic spaces is invariant under quasi-isometry (this is not true for metric spaces in general). In fact $H \leq G$ is quasiconvex if and only if H admits a finite generating set T and the inclusion $H \hookrightarrow X$ is a quasi-isometric embedding with respect to the word metric defined by T .

Given a hyperbolic group G , let

$$\text{Sub}_{\text{QC}[\infty]}(G) \subseteq \text{Sub}_{[\infty]}(G)$$

denote the set of quasiconvex subgroups of G with infinite index. Observe that $\text{Sub}_{\text{QC}[\infty]}(G)$ is invariant under the G -action by conjugation. If G is non-elementary, we prove that this subset is in fact a G -invariant subspace of the perfect kernel of G .

Theorem 6.1. *Let G be a non-elementary hyperbolic group. Then*

$$\overline{\text{Sub}_{\text{QC}[\infty]}(G)} \subseteq \mathcal{K}(G).$$

Recall that a non-elementary hyperbolic group G has a unique maximal finite normal subgroup $N \trianglelefteq G$ and by Proposition 3.2 the set $\text{Conj}(N)^G$ of G -conjugacy classes of subgroups of N defines a finite G -invariant clopen partition: $\text{Sub}(G) = \bigsqcup_{C \in \text{Conj}(N)^G} \{H \leq G : H \cap N \in C\}$. In analogy with Theorem 5.11, we denote for each $C \in \text{Conj}(N)^G$ the (which will be shown to be perfect) subspace

$$\mathcal{F}_C := \overline{\text{Sub}_{\text{QC}[\infty]}(G)} \cap \{H \leq G : H \cap N \in C\} \quad (18)$$

so that the clopen partition of $\text{Sub}(G)$ induces a G -invariant partition of $\overline{\text{Sub}_{\text{QC}[\infty]}(G)}$:

$$\overline{\text{Sub}_{\text{QC}[\infty]}(G)} = \bigsqcup_{C \in \text{Conj}(N)^G} \mathcal{F}_C$$

whose dynamics satisfy:

Theorem 6.2. *Let G be a non-elementary hyperbolic group and fix $C \in \text{Conj}(N)^G$. Then*

- (i) *the action of G on \mathcal{F}_C is topologically transitive, and*
- (ii) *if $|C| = 1$ then the action $G \curvearrowright \mathcal{F}_C$ is highly topologically transitive.*

In particular, if $N = \{\text{id}\}$ then the action $G \curvearrowright \overline{\text{Sub}_{\text{QC}[\infty]}(G)}$ is highly topologically transitive.

Note that the assumption that $|C| = 1$ is a necessary condition for a topologically r -transitive action on \mathcal{F}_C whenever $r \geq 2$ (see Remark 5.13).

We delay the proofs of Theorems 6.1 and 6.2 until the next section. The following is an immediate consequence.

A hyperbolic group is **locally quasiconvex** if every finitely generated subgroup of G is quasiconvex.

Corollary 6.3. *If a group G is non-elementary hyperbolic and locally quasiconvex, then*

$$\mathcal{K}(G) = \text{Sub}_{[\infty]}(G) = \overline{\text{Sub}_{\text{QC}[\infty]}(G)}.$$

If, in addition, the maximal finite normal subgroup of G is trivial then the action of G on $\mathcal{K}(G)$ is highly topologically transitive.

Example 6.4. Hyperbolic limit groups are locally quasiconvex by [Dah03, Proposition 4.6] and torsion free. Therefore, the above corollary provides another proof that the perfect kernel of a hyperbolic limit group $G \neq \mathbb{Z}$ is the space $\text{Sub}_{[\infty]}(G)$ and that the G -action on $\mathcal{K}(G)$ is highly topologically transitive.

Example 6.5. If G is non-elementary hyperbolic and admits a small hierarchy then by [BW13, Theorem 3.6] it is locally quasiconvex and Corollary 6.3 applies.

For instance, if G is not virtually cyclic and splits non-trivially as a graph of groups where each vertex group is a free group, each edge group is cyclic and G does not contain any subgroup which is a Baumslag-Solitar group then G is non-elementary hyperbolic by [BF92] and locally quasiconvex by the above. Compare with the drastically different behaviour when G is a Baumslag-Solitar group [CGLMS22].

Example 6.6. Let S be a finite set with $|S| \geq 2$ and consider the one-relator group $G = \langle S \mid w^m \rangle$ where w is a freely and cyclically reduced word in $S \cup S^{-1}$ and $m \geq |w|_S$. Such one-relator groups with torsion are non-elementary hyperbolic and have no non-trivial normal subgroups [New68, Theorems 3 and 2], and they are locally quasiconvex by [HW01, Theorem 1.2] so, by Corollary 6.3, $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$ and the action $G \curvearrowright \mathcal{K}(G)$ is highly topologically transitive.

Example 6.7. Let G be the Coxeter group $\langle s_1, \dots, s_n \mid s_i^2, (s_i s_j)^{m_{i,j}} \forall i < j \rangle$ where $\infty \geq m_{i,j} > n \geq 3$. Then G is hyperbolic [Mou88, Theorem 17.1] and locally quasiconvex [MW08, Theorem 12.2], so $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$. Moreover, it follows from [PV23] that G has no non-trivial finite normal subgroups so the action $G \curvearrowright \mathcal{K}(G)$ is highly topologically transitive.

Remark 6.8. It is likely that the results of this section can be extended to include relatively hyperbolic groups with suitable parabolic subgroups, for instance finitely generated abelian groups.

6.2 Proof of Theorems 6.1 and 6.2

The proofs of both the main results of this section depend on the fact that, in a non-elementary hyperbolic group G , any infinite index quasiconvex subgroup H is a factor in a splitting of a

quasiconvex subgroup L of G over a finite subgroup (see Proposition 6.11). Moreover, there is a great deal of freedom in the choice of L . In the torsion-free case, this result was first observed by Gromov in [Gro87, 5.3] and was developed in a detailed and effective manner by Arzhantseva in [Arz01]. Other combination theorems allowing torsion were proved by Gitik in [Git99] and were generalised to relatively quasiconvex subgroups of relatively hyperbolic groups by Martínez-Pedroza in [MP09]. In fact, even when restricted to hyperbolic groups, the results of Martínez-Pedroza improve upon Gitik's results in some respects (see for instance Theorems 6.9 and 6.10 below). These improvements are needed for various aspects of our proofs, and they allow us to obtain Proposition 6.11 for hyperbolic groups with torsion.

The Gromov boundary ∂G of a non-elementary hyperbolic group G is a perfect compact metrisable space; it is equipped with a G -action that extends the left multiplication action of G on itself. Given a subgroup H , let $\Lambda(H) \subseteq \partial G$ denote the limit set of H . If H is virtually infinite cyclic then $\Lambda(H)$ consists of precisely two points (the limit points of any infinite order element of H). Let $g \in G$ be an infinite order element and $P = \text{Stab}(\Lambda(\langle g \rangle))$. Then $\langle g \rangle$ is a finite index subgroup of P [GH90, Theorem 30] which implies that P is quasiconvex [Gro87, 8.1D]. Moreover if $a \in G \setminus P$ then $P \cap aPa^{-1} = \text{Stab}(\Lambda(\langle g \rangle) \cap \text{Stab}(\Lambda(\langle ag \rangle)))$ where $\Lambda(\langle g \rangle) \neq \Lambda(\langle ag \rangle)$. It follows that there is a finite index subgroup Q of $P \cap aPa^{-1}$ which fixes three points of ∂G ; this implies that Q fixes a bounded subset of G and is therefore finite. Thus P is **almost malnormal** (i.e. $P \cap P^a$ is finite for all $a \in G \setminus P$; here and in the sequel, P^a denotes the conjugate aPa^{-1}). By [Bow12, Theorem 7.11] this implies that G is hyperbolic relative to P . Since subgroups of conjugates of P are virtually cyclic and therefore quasiconvex it follows from [MP12, Theorem 1.1] that quasiconvexity in G is equivalent to quasiconvexity relative to P . In particular, the following theorems are special cases of the results of [MP09].

Theorem 6.9 (Martínez-Pedroza, [MP09, Theorem 1.1]). *Let G be a hyperbolic group and fix a finite generating set S for G . Let $g \in G$ be an infinite order element and $P = \text{Stab}(\Lambda(\langle g \rangle))$. Let $H \leq G$ be a quasiconvex subgroup. Then there is a constant $C_1 \geq 0$ depending only on H, P and S such that the following holds. If $Q \leq P$ is a subgroup such that*

1. $H \cap P \leq Q$ and
2. $|k|_S \geq C_1$ for all $k \in Q \setminus H$

*then $\langle H, Q \rangle$ splits as the amalgamated free product $\langle H, Q \rangle = H *__{H \cap Q} Q$ and $\langle H, Q \rangle$ is quasiconvex in G .*

Theorem 6.10 (Martínez-Pedroza, [MP09, Theorem 1.2]). *Let G be a hyperbolic group and fix a finite generating set S for G . Let $g \in G$ be an infinite order element and $P = \text{Stab}(\Lambda(\langle g \rangle))$. Let $H_1, H_2 \leq G$ be quasiconvex subgroups such that $M := H_1 \cap P = H_2 \cap P$. Then there exists a constant $C_2 \geq 0$ depending only on H_1, H_2, P and S such that the following holds. If $p \in P$ is such that*

1. $pMp^{-1} = M$
2. $|k|_S \geq C_2$ for all $k \in MpM$

*then $\langle H_1, pH_2p^{-1} \rangle$ splits as the amalgamated free product $\langle H_1, pH_2p^{-1} \rangle = H_1 *_M pH_2p^{-1}$ and $\langle H_1, pH_2p^{-1} \rangle$ is quasiconvex in G .*

We are now in a position to prove the combination result we need. In order to produce the relevant subgroup $P \leq G$, we adapt a standard argument to produce a suitable hyperbolic element of G by taking advantage of the “holes” in the boundary of G left by the subgroups H_i .

Proposition 6.11. *Let G be a non-elementary hyperbolic group and $H_1, \dots, H_r \leq G$ be quasiconvex subgroups with infinite index. Let $N \trianglelefteq G$ be the unique maximal finite normal subgroup of G and suppose that $M := H_1 \cap N = H_i \cap N$ for each i . Then the following hold.*

1. *For each i , the subset $\Lambda(H_i) \subseteq \partial G$ is closed with empty interior. Thus the complement $\partial G \setminus \cup_{i=1}^r \Lambda(H_i)$ is a dense open set.*
2. *For any non-empty open set $Y \subseteq \partial G$, there exists an infinite order element $p \in G$ whose limit points are contained in Y and such that, if we denote $P = \text{Stab}(\Lambda(\langle p \rangle))$, then $P \cap H_i = M$ for each i .*
3. *If $u \in G$ is a sufficiently high power of p then, for each $n \in \mathbb{N}$, the subgroup $Q_n := \langle u^n, M \rangle$ generated by u^n and M splits as the direct product $Q_n = \langle u^n \rangle \times M$, and for each $i \in \{1, \dots, r\}$, the subgroup $\langle H_i, Q_n \rangle$ splits as the amalgamated free product*

$$\langle H_i, Q_n \rangle = H_i *_M Q_n$$

and is quasiconvex and has infinite index in G .

Proof. Recall that the unique maximal finite normal subgroup $N \trianglelefteq G$ is the kernel of the action of G on ∂G (see e.g. [DK18, Lemma 11.130]). Let $H \leq G$ be an infinite index quasiconvex subgroup. Then the limit set $\Lambda(H)$ is a (possibly empty, if H is finite) closed proper subspace of ∂G . Suppose for a contradiction that there is an open subset $U \subseteq \partial G$ such that $U \subseteq \Lambda(H)$. Then by [Gro87, 8.2G], there is an infinite order element $h \in H$ such that $\Lambda(\langle h \rangle) \subseteq U$. Since h acts on ∂G with north-south dynamics ([Gro87, 8.1G]), a certain power h^n satisfies $h^n U \cup U = \partial G$. But $h^n U \cup U \subseteq h^n \Lambda(H) \cup \Lambda(H) = \Lambda(H)$. Thus $\Lambda(H) = \partial G$ which contradicts the fact that H has infinite index in G . Thus $\Lambda(H)$ has empty interior. The same holds for a finite union of closed sets with empty interior. This proves part 1 of the lemma.

To prove part 2, we first show that, for any $g \in G \setminus N$, the fixed point set $\text{Fix}(g) := \{x \in \partial G : gx = x\}$ is closed with empty interior. Fix an element $g \in G$. First, $\text{Fix}(g)$ is closed. Suppose that there is a non-empty open set $W \subseteq \text{Fix}(g)$ and let $L \leq G$ be the pointwise stabiliser of W . Since ∂G is perfect, W contains more than two points so every element of L has finite order. By [GH90, Corollaire 36, Chapitre 8] this implies that L is finite. Let $h \in G$ be an infinite order element whose limit points h^+, h^- are in W and let $W^+, W^- \subseteq W$ be open neighbourhoods of h^+, h^- respectively such that $W^+ \cap W^- = \emptyset$. Let L^+ be the pointwise stabiliser of W^+ and note that, as above, L^+ is finite. Moreover, for sufficiently large $n \in \mathbb{N}$, $h^n W^- \cup W^+ = \partial G$ and in particular $W^- \subseteq h^n W^-$, so $h^n L^+ h^{-n} \leq L^+$. Since L^+ is finite, this implies that $h^n L^+ h^{-n} = L^+$. But then g is in the pointwise stabiliser of $h^n W^-$ and of W^+ so $g \in N$.

Recall that any subgroup $H \leq G$ acts properly discontinuously on $\partial G \setminus \Lambda(H)$ ([Coo89]). Let $Y \subseteq \partial G$ be non-empty and open and let $x_0 \in Y' := Y \cap (\partial G \setminus \cup_{i=1}^r \Lambda(H_i))$. Then for each i there is an open neighbourhood $U_i \subseteq Y'$ such that $|\{h \in H_i : hU_i \cap U_i \neq \emptyset\}| < \infty$. Let $U := \cap_{i=1}^r U_i$ and note that U is non-empty and open. Let $A = \{h \in (\cup_{i=1}^r H_i) \setminus M :$

$hU \cap U \neq \emptyset$. Since, for each $h \in A$, the fixed point set $\text{Fix}(h)$ is closed with empty interior, the complement $V := U \setminus (\cup_{h \in A} \text{Fix}(h))$ is non-empty and open. Now enumerate $A = \{a_1, \dots, a_l\}$. If $a_1 V \cap V = \emptyset$, let $V_1 := V$. Otherwise there exists $x_1 \in V$ such that $a_1 x_1 \in V$ and, since $x_1 \notin \text{Fix}(a_1)$, we have $x_1 \neq a_1 x_1$. It follows that there exists an open neighbourhood $V_1 \subseteq V$ of x_1 such that $a_1 V_1 \cap V_1 = \emptyset$. Iterate this process to find a sequence of non-empty open sets $V_l \subseteq \dots \subseteq V_1 \subseteq V$ such that for each $1 \leq k \leq l$ and each $1 \leq i \leq k$ we have $a_i V_k \cap V_k = \emptyset$. In particular $a V_l \cap V_l = \emptyset$ for all $a \in A$.

Let $p \in G$ be an infinite order element whose limit points are contained in $V_l \subseteq Y$ and let $P = \text{Stab}(\Lambda(\langle p \rangle))$. Then $P \cap H_i = M$ for each i . This proves part 2 of the lemma.

Let u be a power of p which commutes with M . For each $n \in \mathbb{N}$ let $Q_n := \langle u^n, M \rangle = \langle u^n \rangle \times M \leq P$. Then $Q_n \cap H_i = M$ for each i and, since P is quasi-isometrically embedded in G , for sufficiently large n and $C_1 \geq 0$ as in Theorem 6.9, $|k|_S \geq C_1$ for all $k \in Q_n \setminus M$. Therefore, by Theorem 6.9, for sufficiently large n ,

$$\langle H_i, Q_n \rangle = H_i *_M Q_n \leq G \quad (19)$$

is quasiconvex. Moreover, up to replacing u with u^{n_0} for some sufficiently large n_0 , each $\langle H_i, Q_n \rangle$ is an infinite index quasiconvex subgroup of G satisfying (19). Indeed, if $H_i *_M Q_{n_0}$ has finite index, then the splitting is non-trivial (if it were trivial, i.e. $H_i = M$, and G would be elementary), thus $\langle H_i, Q_{2n_0} \rangle$ has infinite index in $\langle H_i, Q_{n_0} \rangle$ and therefore in G . This proves part 3 of the lemma. \square

Proof of Theorem 6.1. Let $H \leq G$ be an infinite index quasiconvex subgroup, $N \trianglelefteq G$ be the unique maximal finite normal subgroup of G and $M = H \cap N$. By Proposition 6.11, there exists an infinite order element $u \in G$ such that, for each $n \in \mathbb{N}$, $Q_n := \langle u^n, M \rangle = \langle u^n \rangle \times M$ and

$$H_n := \langle H, Q_n \rangle = H *_M Q_n \in \text{Sub}_{QC[\infty]}(G).$$

Let T be a finite generating set for H and $\lambda \geq 1$ be such that the natural inclusion of the first term of the sequence $H_1 \hookrightarrow G$ is a λ -quasi-isometric embedding with respect to the generating set S for G and the generating set $T \cup \{u\}$ for H_1 . For any $n \in \mathbb{N}$, if $o \in H_n \setminus H$ then the reduced spelling of o must contain the subword u^n , and since $o \in H_1 \setminus H$ we have

$$|o|_S \geq \frac{1}{\lambda} |o|_{T \cup \{u\}} - \lambda \geq \frac{n}{\lambda} - \lambda.$$

Hence, given a basic open neighbourhood $\mathcal{V}(\mathcal{I}, \mathcal{O})$ of H in $\text{Sub}(G)$, for all sufficiently large n , the subgroup H_n contains H but does not contain $o \in \mathcal{O}$, thus $H_n \in \mathcal{V}(\mathcal{I}, \mathcal{O}) \cap \text{Sub}_{QC[\infty]}(G)$ so H is not isolated in $\text{Sub}_{QC[\infty]}(G)$. \square

Proof of Theorem 6.2. Fix $C \in \text{Conj}(N)^G$. To prove part (i) (i.e. that the action of G on \mathcal{F}_C is topologically transitive) we will show the following. Given two non-empty open subsets $U, V \subseteq \mathcal{F}_C$, there exists $g \in G$ such that $(g \cdot U) \cap V = gUg^{-1} \cap V \neq \emptyset$. As in the proof of Theorem 5.11, we can assume, up to replacing V with its image by some element of G , that there exist finitely generated subgroups $H \in U$ and $K \in V$ such that $M := H \cap N = K \cap N$.

To prove part (ii), we need to show that, if $C = \{M\}$ (i.e. M is normal in G) then, for all r and for all non-empty open sets $U_1, \dots, U_r, V_1, \dots, V_r \subseteq \mathcal{F}_C$, there exists $g \in G$ such that $gU_i \cap V_i \neq \emptyset$ for each i . Since finitely generated subgroups are dense and, for all $H_i \in U_i$ and

$K_i \in V_i$, we have $H_i \cap N = K_i \cap N = M$, both part (i) and part (ii) follow from the following statement:

Fix $C \in \text{Conj}(N)^C$ and $M \in C$ (we don't assume that $|C| = 1$). Let $r \geq 1$ and $U_1, \dots, U_r, V_1, \dots, V_r \subseteq \mathcal{F}_C$ be open subsets such that, for each $i \in \{1, \dots, r\}$, there exist finitely generated subgroups $H_i \in U_i$ and $K_i \in V_i$ where $H_i \cap N = K_i \cap N = M$. Then there is a common element $g \in G$ such that $gU_i \cap V_i \neq \emptyset$ for all i .

Let us prove that this statement indeed holds.

Claim 1. *There exists an infinite order element $p \in G$ which commutes with M and such that, for each i , $\langle H_i, K_i^p \rangle = H_i *_M K_i^p \leq G$ is quasiconvex and has infinite index in G .*

Proof of Claim 1. If G has infinitely many ends, then it admits a Stallings tree, a minimal irreducible G -tree T with finite edge stabilisers. Note that the kernel of the action on T is the maximal finite normal subgroup N . The infinite index subgroups H_i, K_i have proper minimal subtrees T_{H_i}, T_{K_i} in T ; let Z_e be the half-tree separated from the union $\cup_i T_{H_i} \cup \cup_i T_{K_i}$ of these minimal subtrees by some edge e such that $\text{Stab}_{H_i}(e) = \text{Stab}_{K_i}(e) = M$ for each i (see Claim 4 in the proof of Theorem 5.11). Take a vertex x of T . The map $G \ni g \mapsto g \cdot x$ extends to a G -equivariant map $\Psi : G \cup \partial G \rightarrow T \cup \partial T$. By Lemma 5.2 there is a hyperbolic (with respect to the action on T) element $p \in G$ whose axis $\text{Axis}(p)$ is contained in Z_e . Since it is fixed by $\langle p \rangle$, the image $\Psi(\Lambda(\langle p \rangle))$ is the set of limit points of $\text{Axis}(p)$. Thus, the stabiliser $P := \text{Stab}_G(\Lambda(\langle p \rangle))$ stabilises globally $\text{Axis}(p)$. It follows that any $g \in P \cap H_i$ (respectively of $P \cap K_i$), for any i , must stabilise $\text{Axis}(p)$ and T_{H_i} (respectively T_{K_i}), therefore $g \in \text{Stab}_{H_i}(e) = \text{Stab}_{K_i}(e) = M$.

Otherwise, if G has only one end, let Y any open subset of ∂G . By Proposition 6.11 there exists an infinite order element $p \in G$ whose limit points are contained in Y such that, if $P := \text{Stab}(\Lambda(\langle p \rangle))$, then $P \cap H_i = P \cap K_i = M$ for each i .

In either case, replace p with a proper power so that it commutes with M .

Now let $C_2 \geq 0$ be as in Theorem 6.10. Since $\langle p \rangle$ is quasi-isometrically embedded in G and M is finite, we can replace p with a proper power so that $|pm| \geq |p| - |m| \geq C_2$ for all $pm \in pM = MpM$. Thus by Theorem 6.10, for each i , the subgroup generated by H_i and K_i^p splits as

$$\langle H_i, K_i^p \rangle = H_i *_M K_i^p,$$

and is quasiconvex in G .

Moreover, up to taking a proper power of p , we can assume that for each i , $H_i *_M K_i^p$ has infinite index in G . Indeed, suppose that $H_i *_M K_i^p$ has finite index in G for some i . Then the splitting is non-trivial (if M is equal to one of the factors then the other factor has finite index in G) so G has at least two, thus infinitely many, ends. In this case, p acts hyperbolically on the Stallings tree with axis $\text{Axis}(p)$ contained in Z_e . Let $n \geq 2$ and let $\pi(e)$ be the projection of e on $\text{Axis}(p)$. Then any edge $e' \not\subset \text{Axis}(p)$ but adjacent to $\text{Axis}(p)$ at a vertex strictly between $\pi(e)$ and $p^n \cdot \pi(e)$ defines a half-tree Z' (the one that does not contain e) which lies outside any minimal invariant $H_i *_M K_i^{p^n}$ -subtree of T . Thus the quotient $H_i *_M K_i^{p^n} \backslash T$ is infinite. Since the action of G on T is minimal and G is finitely generated, the quotient $G \backslash T$ is compact (see Remark 5.6) so this implies that $H_i *_M K_i^{p^n}$ has infinite index in G . \diamond

By Proposition 6.11-(3), there exists an infinite order element $u \in G$ which commutes

with M and such that, for all i ,

$$\begin{aligned} A^{(i)} &:= \langle H_i, K_i^p, u \rangle = (H_i *_M K_i^p) *_M \langle u, M \rangle \leq G \quad \text{and} \\ B^{(i)} &:= (A^{(i)})^{p^{-1}} = \langle H_i^{p^{-1}}, K_i, u^{p^{-1}} \rangle = (H_i^{p^{-1}} *_M K_i) *_M \langle u^{p^{-1}}, M \rangle \leq G \end{aligned}$$

are quasiconvex and have infinite index in G . For ease of notation, we now fix $i \in \{1, \dots, r\}$ and denote $A = A^{(i)}$ and $B = B^{(i)}$.

Claim 2. *For all large enough n , we have $A_n := H_i *_M (K_i)^{p^{u^n}} \in U_i$ while $B_n = (A_n)^{u^{-n}p^{-1}} = (H_i)^{u^{-n}p^{-1}} *_M K_i \in V_i$.*

Proof of Claim 2. First observe, for each n , that $A_n \leq A$, $B_n \leq B$ and the inclusions $A_n \hookrightarrow A$ and $B_n \hookrightarrow B$ are quasi-isometric embeddings so A_n, B_n are quasiconvex in G and have infinite index. Moreover $A_n \cap N = B_n \cap N = M$ so $A_n, B_n \in \mathcal{F}_C$. Now let $\mathcal{V}(\mathcal{I}, \mathcal{O}) \subseteq \text{Sub}(G)$ be a basic neighbourhood of H_i such that $\mathcal{V}(\mathcal{I}, \mathcal{O}) \cap \mathcal{F}_C \subseteq U_i$. Let T, T' be finite generating sets for H_i, K_i respectively and let $\lambda \geq 1$ be such that $A \hookrightarrow G$ is a λ -quasi-isometry with respect to the word metric on G defined by S and the word metric on A defined by $S_A := T \cup pT'p^{-1} \cup \{u\}$. Then for all $a \in A_n \setminus H_i$, the reduced spelling of a must contain a subword $pu^n t' u^{-n} p^{-1}$, thus

$$|a|_S \geq \frac{1}{\lambda} |a|_{S_A} - \lambda \geq \frac{2n}{\lambda} - \lambda.$$

Therefore if n is sufficiently large then $|a|_S > |o|_S$ for all $a \in A_n \setminus H_i$ and $o \in \mathcal{O}$. Since $\mathcal{I} \subseteq H_i \subseteq A_n$, this implies that $A_n \in \mathcal{V}(\mathcal{I}, \mathcal{O}) \cap \mathcal{F}_C \subseteq U_i$. By a symmetric argument, $(u^{-n}p^{-1}) \cdot A_n = B_n \in V_i$ for sufficiently large n . \diamond

Hence, by applying Claim 2 to each $i \in \{1, \dots, r\}$, we find $N \in \mathbb{N}$ such that, for each i , the intersection $(u^{-N}p^{-1} \cdot U_i) \cap V_i$ is non-empty. \square

6.3 Hyperbolic 3-manifold groups

In the case of hyperbolic 3-manifold groups the results of the previous section, together with celebrated results about 3-manifolds, allow us to describe the perfect kernel completely.

The following dichotomy, which is a consequence of Canary's Covering Theorem [Can96] and the Tameness Theorem [Ago04, CG06] (see [AFW15, Theorem 5.2] for a precise statement) allows us to understand the status of non-quasiconvex subgroups in the space of subgroups of a hyperbolic 3-manifold groups. If $G = \pi_1(M)$, where M is a hyperbolic 3-manifold, and $H \leq G$ is finitely generated then one (and only one) of the following holds:

- (a) H is quasiconvex or
- (b) H is a **virtual fiber**, i.e. there exists a finite index subgroup $G' \leq G$ such that $H' := G' \cap H$ is a hyperbolic surface group and G' fits in a short exact sequence

$$1 \longrightarrow H' \longrightarrow G' \longrightarrow \mathbb{Z} \longrightarrow 1.$$

Theorem 6.12. *Let M be a closed hyperbolic 3-manifold and $G = \pi_1(M)$. Then*

$$\mathcal{K}(G) = \overline{\text{Sub}_{\text{QC}[\infty]}(G)}.$$

More precisely, the perfect kernel of G is the set of infinite index subgroups $H \leq G$ such that either H is quasiconvex or H is not finitely generated.

Moreover, the action of G on $\mathcal{K}(G)$ is highly topologically transitive and the Cantor-Bendixson rank of G is either 2 or 3.

Proof. By Theorem 6.1 and since G is finitely generated $\overline{\text{Sub}_{\text{QC}[\infty]}(G)} \subseteq \mathcal{K}(G) \subseteq \text{Sub}_{[\infty]}(G)$. Let us show that the first inclusion is in fact an equality.

If $H \leq G$ is a virtual fiber subgroup then, by Corollary 3.6, $H \notin \mathcal{K}(G)$.

It remains to show that every non-finitely generated subgroup of G is the limit of a sequence of quasiconvex subgroups with infinite index.

It follows from Proposition 3.15 that a non-finitely generated subgroup does not contain any virtual fiber subgroup. Indeed, for any tower of subgroups $K_1 \leq K_2 \leq G$ where K_1 is a virtual fiber subgroup, either $[K_1 : K_2] < \infty$ or $[K_2 : G] < \infty$ and in either case K_2 is finitely generated. Thus if $H = \langle s_n : n \in \mathbb{N} \rangle \leq G$ is not finitely generated, it is the limit of the non-stationary sequence $(\langle s_1, \dots, s_n \rangle)_n$ of infinite index subgroups and each $\langle s_1, \dots, s_n \rangle$ is quasiconvex.

To see that $\text{rk}_{\text{CB}}(G) = 2$ or 3, first note that the Cantor-Bendixson rank of G is the supremum of the Cantor-Bendixson erasing ranks of its finite index and virtual fiber subgroups. But virtual fiber subgroups are separable. Indeed if $G'/H' = \mathbb{Z}$ then $H' = \bigcap_{n \in \mathbb{N}} \pi^{-1}(n\mathbb{Z})$ where $\pi : G' \rightarrow \mathbb{Z}$ is the quotient map. For each $n \in \mathbb{N}$, $\pi^{-1}(n\mathbb{Z})$ has finite index in G' and therefore in G , so H' is closed with respect to the profinite topology on G . Since H is the union of finitely many cosets of H' , it follows that H is also closed. Thus $\text{rk}_{\text{CB}}^\times(H; G) \geq 2$. Since by [Ago13, Theorem 9.2] every closed hyperbolic 3-manifold group contains a virtual fiber subgroup, this implies that $\text{rk}_{\text{CB}}(G)$ is in fact the supremum of the Cantor-Bendixson erasing ranks of the virtual fiber subgroups. For any such subgroup $H \leq G$, any Schreier graph of H is quasi-isometric to \mathbb{Z} so it follows from Theorem 3.10 that $\text{rk}_{\text{CB}}^\times(H; G) \leq 3$. \square

Note that there exist hyperbolic 3-manifold groups with Cantor-Bendixson rank 3. Indeed, it was pointed out to us by Ian Biringer that there are hyperbolic 3-manifold groups which algebraically fiber over the infinite dihedral group D_∞ and the kernel of this fibering has Cantor-Bendixson erasing rank 3. Recall that a group G **algebraically fibers** over Q if it fits into a short exact sequence $1 \rightarrow H \rightarrow G \rightarrow Q \rightarrow 1$ where H is finitely generated. He moreover convinced us (personal communication) that if a virtual fiber subgroup is not virtually of this form then its Cantor-Bendixson erasing rank is 2.

It follows that the Cantor-Bendixson rank of a hyperbolic 3-manifold group G is 3 if and only if G virtually fibers over D_∞ , otherwise it is 2. This leads us to the following question.

Question 6.13. Does there exist a closed hyperbolic 3-manifold M such that no finite index subgroup of $\pi_1(M)$ algebraically fibers over D_∞ ? Equivalently, does there exist a hyperbolic 3-manifold group G with $\text{rk}_{\text{CB}}(G) = 2$?

7 Some facts and more questions

Let us consider the supremum of the Cantor-Bendixson ranks among various classes of groups

$$\begin{aligned}\alpha_{\text{fp}} &:= \sup\{\text{rk}_{\text{CB}}(G) \mid G \text{ finitely presented}\} \\ \alpha_{\text{fg}} &:= \sup\{\text{rk}_{\text{CB}}(G) \mid G \text{ finitely generated}\} \\ \alpha_c &:= \sup\{\text{rk}_{\text{CB}}(G) \mid G \text{ countable}\}\end{aligned}$$

Question 7.1. Clearly $\alpha_{\text{fp}} \leq \alpha_{\text{fg}} \leq \alpha_c$.

1. Which inequalities are strict?
2. What are the values of α_{fp} , α_{fg} and α_c ?
3. What are the possible values of Cantor-Bendixson ranks for the various classes of groups?

Recall the **Rips construction** and the **Haglund-Wise construction**:

Given any finitely presented group Q there exists a specific group G and a finitely generated normal subgroup N with a short exact sequence

$$1 \longrightarrow N \longrightarrow G \xrightarrow{\phi} Q \longrightarrow 1,$$

where the group G can be chosen to be

- a hyperbolic group [Rip82], or
- a finitely presented subgroup of $\text{SL}(n, \mathbb{Z})$ for some n [HW08, Th. 1.6].

It follows that:

Proposition 7.2.

$$\begin{aligned}\alpha_{\text{fp}} &= \sup\{\text{rk}_{\text{CB}}(G) \mid G \text{ hyperbolic}\}. \\ \alpha_{\text{fp}} &= \sup\{\text{rk}_{\text{CB}}(G) \mid G \text{ finitely presented linear}\}.\end{aligned}$$

Recall from Proposition 3.5 that given an exact sequence $1 \rightarrow N \rightarrow G \xrightarrow{\phi} Q \rightarrow 1$ where N is finitely generated, then $\text{Sub}_{N \leq}(G)$ is a G -invariant clopen subset of $\text{Sub}(G)$ and ϕ induces a homeomorphism $\text{Sub}_{N \leq}(G) \simeq \text{Sub}(Q)$ that is G -equivariant for the G -actions, where G acts on $\text{Sub}(Q)$ via ϕ , so that $\mathcal{K}(G) \cap \text{Sub}_{N \leq}(G) \simeq \mathcal{K}(Q)$ and $\text{rk}(G) \geq \text{rk}(Q)$ (Proposition 3.5 (4)).

It follows from the Rips and Haglund-Wise constructions that for any finitely presented Q we can embed the action $Q \curvearrowright \text{Sub}_{[\infty]}(Q)$, regardless of its complexity, inside the space of non quasiconvex subgroups (see Corollary 7.4) of some hyperbolic group, or alternatively inside the space of subgroups of a finitely presented linear group.

Question 7.3. These observations give rise to the questions:

1. How “pathological” can the action $H \curvearrowright \text{Sub}(H)$ be for a finitely presented group H ?
2. Which bounds can be set for the Cantor-Bendixson ranks of finitely presented groups?

Observe that the lamplighter groups, the Grigorchuk group and the Gupta-Sidki 3-group mentioned in the introduction are not finitely presented. The first example of a finitely presented group with infinite Cantor-Bendixson rank (namely the Baumslag group) has been announced by Bontemps, Frisch and the second author [BFG23]. This implies in particular the existence of hyperbolic groups with infinite Cantor-Bendixson rank.

Corollary 7.4. *If G is produced using the Rips construction above with a non-hyperbolic Q , then*

$$\text{Sub}_{N \leq}(G) \cap \overline{\text{Sub}_{\text{QC}[\infty]}(G)} = \emptyset.$$

Thus, if $\mathcal{K}(Q) \neq \emptyset$, the reverse inclusion $\overline{\text{Sub}_{\text{QC}[\infty]}(G)} \supseteq \mathcal{K}(G)$ of Theorem 6.1 does not hold since

$$\mathcal{K}(Q) \simeq \mathcal{K}(G) \cap \text{Sub}_{\leq N}(G) \subseteq \mathcal{K}(G) \setminus \overline{\text{Sub}_{\text{QC}[\infty]}(G)}.$$

The corollary follows from the following result of Mihalik and Towle [MT94, Theorem 1]: *If a subgroup of a hyperbolic group G has infinite index and is quasiconvex, then it does not contain any infinite normal subgroup of G .* The fact that Q is not hyperbolic ensures that N is infinite.

The Theorems 5.9, 5.11, 6.1, 6.2 all use the property that the subgroups we consider fix a proper quasiconvex subset of an ambient hyperbolic space. This leads to the following question, which we have not pursued in depth:

Question 7.5. Is there a common generalisation of our results in the context of group actions on hyperbolic spaces with suitable constraints?

8 An application: amenable faithful transitive actions

The class \mathcal{A} is the class of countably infinite groups that admit a faithful, transitive and amenable action.

Recall from [GM07] that if a triple of groups $K_1 \leq K_2 \leq K_3$ satisfies that K_i is co-amenable in K_{i+1} , then K_1 is co-amenable in K_3 and if K_1 belongs to \mathcal{A} , then K_3 belongs to \mathcal{A} . In particular, if $H \in \mathcal{A}$ and $G \geq H$ is a finite index supergroup, then $G \in \mathcal{A}$. Moreover, if $H_1, H_2 \in \mathcal{A}$ then $H_1 \times H_2 \in \mathcal{A}$.

Moon observed that Baumslag-Solitar groups belong \mathcal{A} since they admit a co-amenable free subgroup (or they are amenable, depending on the parameters).

Proposition 8.1. *Let G be a countable group and $\text{Sub}_{[\infty]}(G)$ its space of infinite index subgroups. Assume that there is*

- (a) *a subgroup $H_1 \in \text{Sub}_{[\infty]}(G)$ such that $G \curvearrowright G/H_1$ is faithful;*
- (b) *a co-amenable subgroup $H_2 \in \text{Sub}_{[\infty]}(G)$ (or more generally a family $(H_2^i)_{i \in \mathbb{N}}$ of subgroups $H_2^i \in \text{Sub}_{[\infty]}(G)$ such that the action $G \curvearrowright G/H_2^1 \sqcup G/H_2^2 \sqcup \dots \sqcup G/H_2^i \sqcup \dots$ is amenable);*
- (c) *a subgroup $H_0 \in \text{Sub}_{[\infty]}(G)$ whose orbit under the G -action by conjugation on $\text{Sub}_{[\infty]}(G)$ accumulates on both H_1 and H_2 (or more generally on H_1 and all the H_2^i 's).*

Then the action $G \curvearrowright G/H_0$ is faithful, transitive and amenable. In particular, $G \in \mathcal{A}$.

Moreover, if $R \subseteq \text{Sub}(G)$ is a G -invariant closed subspace containing both H_1 and H_2 (resp. H_1 and all the H_2^i) together with a dense orbit, then the set of H_0 satisfying (c) is a dense G_δ subset of R .

Proof of Corollary I. First observe that the group G is infinite, thus $H_1 := \{\text{id}\} \in \text{Sub}_{[\infty]}(G)$. The amenability condition on G implies that it admits subgroups satisfying Condition (b) of Proposition 8.1, and thus its conclusion. \square

Proof of Proposition 8.1. It is enough to show that $G \curvearrowright G/H_0$ is faithful and amenable. Let $\gamma \in G \setminus \{\text{id}\}$. By faithfulness of $G \curvearrowright G/H_1$, there is $g_1 \in G$ such that $g_1^{-1}\gamma g_1 \notin H_1$. Let R be the closure of the orbit of H_0 and let V be the open neighbourhood of H_1 in R consisting of the subgroups that do not contain $g_1^{-1}\gamma g_1$. By density, there is $g \in G$ such that $gH_0g^{-1} \in V$, i.e. $\gamma \notin g_1gH_0(g_1g)^{-1}$. Thus $G \curvearrowright G/H_0$ is faithful.

Consider a Følner sequence $(A_i)_i$ with $A_i \subseteq G/H_2^i$ (or $A_i \subseteq G/H_2$). More precisely, one can (up to re-indexing) write G as an increasing union of finite subsets $G = \bigcup_i S_i$ such that for each i and $\gamma \in S_i$ we have $|\gamma A_i \Delta A_i| \leq \frac{1}{i}|A_i|$ with $A_i = \{g_j^i H_2^i : j = 1, 2, \dots, |A_i|\}$. Let $|A_i| = n_i$ and $J_i = \{1, 2, \dots, n_i\}$. The intersection $\gamma A_i \cap A_i$ is encoded by the equations $(g_k^i)^{-1}\gamma g_j^i \in H_2^i$ (for $j, k \in J_i$ and $\gamma \in S_i$) while $|A_i|$ is encoded by $(g_k^i)^{-1}\gamma g_j^i \notin H_2^i$ (for $j, k \in J_i, j \neq k$). These conditions define an open neighbourhood V_i of H_2^i in R and by density of the orbit of H_0 , there is $k_i \in G$ such that $k_i H_0 k_i^{-1} \in V_i$, i.e. $(g_k^i)^{-1}\gamma g_j^i \in H_2^i \implies (g_k^i)^{-1}\gamma g_j^i \in k_i H_0 k_i^{-1}$ and $(g_k^i)^{-1}\gamma g_j^i \notin k_i H_0 k_i^{-1}$. It follows that the $B_i := \{g_j^i k_i H_0 : j \in J_i\}$ satisfy $|\gamma B_i \Delta B_i| \leq \frac{1}{i}|B_i|$. Thus the sequence $(B_i)_i$ is a Følner sequence for the action $G \curvearrowright G/H_0$, which is therefore amenable.

Concerning the ‘‘moreover’’ part, since R admits a countable basis $(U_\ell)_{\ell \in \mathbb{N}}$ of open sets, the set $\cap_\ell G \cdot U_\ell$ of points with dense orbits in R is a G_δ . Since their orbit accumulates to some infinite index subgroup (namely H_1), they have infinite index themselves. \square

Corollary 8.2. *Any finite graph of free groups with cyclic edges with at least one non-cyclic vertex group is in \mathcal{A} .*

This generalises a result of Moon [Moo11a].

Proof. We will apply Proposition 8.1. By Example 5.22, G belongs to \mathcal{Q}' . Thus, $G \curvearrowright \text{Sub}_{[\infty]}(G) =: R \ni \{\text{id}\}$ is topologically transitive. It remains to find a co-amenable infinite index subgroup $H_2 \in R$.

If the graph is not a tree, then G surjects onto \mathbb{Z} . Take H_2 to be the kernel of this morphism. Otherwise, if one of the leaves v of the finite graph has a non-cyclic stabiliser G_v , then G surjects onto the abelianisation of G_v modulo the group G_e of the incident edge e . Eventually, if v_1, v_2 are two leaves of the finite graph, with cyclic stabilisers G_{v_1}, G_{v_2} (and the stabilisers $G_{e_1} \not\leq G_{v_1}$ and $G_{e_2} \not\leq G_{v_2}$ of the incident edges e_1, e_2 are proper subgroups). Then G maps onto the non-trivial free product $G_{v_1}/G_{e_1} * G_{v_2}/G_{e_2}$, that itself maps virtually onto \mathbb{Z} . Thus G virtually maps onto \mathbb{Z} . The kernel of this virtual morphism is co-amenable and has infinite index. \square

Corollary 8.3. *Let G be a countable group which admits a minimal and irreducible action $G \curvearrowright T$ on a tree. Suppose that there are two edges f_1, f_2 of T with $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2) = N_0$ (the kernel of the G -action on T), and assume moreover that G admits a co-amenable subgroup G_0 (possibly a finite index subgroup) such that $G_0 \cap N_0 = \{\text{id}\}$ and such that the graph $G_0 \backslash T$ is not a finite tree. Then $G \in \mathcal{A}$.*

Proof of Corollary 8.3. If $G_0 \backslash T$ is not a tree, G_0 admits a homomorphism onto \mathbb{Z} whose kernel N gives rise to a infinite quotient $N \backslash T$: indeed, pick a simple closed loop c in the graph $G_0 \backslash T$, then the fundamental group G_0 of the graph of groups associated with $G_0 \curvearrowright T$ maps onto $\pi_1(c) \simeq \mathbb{Z}$ with kernel $N \leq G_0$. The graph of groups associated with $N \curvearrowright T$ contains an embedded copy of the universal cover of the loop c . It follows that $H_2 := N$

is co-amenable in G_0 and thus in G and that $H_2 \in \overline{\text{Sub}_{|\bullet|_T|_\infty}(G)}$ and $H_2 \cap N_0 = \{\text{id}\}$ (i.e. $H_2 \in \mathcal{F}_{\{\{\text{id}\}\}}^T$ in the terminology of Equation 13 and Theorem 5.11). In case $G_0 \setminus T$ is infinite, then simply set $H_2 := G_0$. By Theorem 5.11, $R := \mathcal{F}_{\{\{\text{id}\}\}}^T$ and H_2 satisfy the assumptions of Proposition 8.1. \square

Corollary 8.4. *Let G be a group which decomposes non-trivially as the fundamental group of a graph of groups where the graph is not a finite tree. Assume that, for some edge e and incident vertex v , the associated edge group G_e and vertex group G_v satisfy: $\exists g \in G_v$ such that $gG_e g^{-1} \cap G_e = \{\text{id}\}$. Then $G \in \mathcal{A}$.*

Proof. The condition $gG_e g^{-1} \cap G_e = \{\text{id}\}$ ensures that in the associated Bass-Serre tree T , the edge f_1 stabilised by $G_e \leq G$ and its image by $g \in G_v$ have stabilisers that intersect trivially. Thus the kernel N_0 of the action is trivial. The corollary follows by applying Corollary 8.3 with $G_0 := G$. \square

Our final application is to show that the class of (virtually) **compact special groups** introduced by Haglund and Wise in [HW08] is contained in \mathcal{A} . These are groups G which (virtually) act co-specially, freely and cocompactly on a CAT(0) cube complex X : a cell complex constructed by attaching n -cubes $[0, 1]^n$ isometrically along faces, which is CAT(0) when equipped with the piecewise ℓ^2 -metric. The action $G \curvearrowright X$ is **co-special** if the quotient $G \backslash X$ is special in the sense of [Wis21, Chapter 6, p. 167].

We recall some key definitions and properties related to CAT(0) cube complexes and co-special actions.

Let X be a CAT(0) cube complex. The **dimension** of X is the supremum of the set of $n \in \mathbb{N}$ such that X contains an n -cube. We say that X is **reducible** if it splits non-trivially as a product $X = X_1 \times X_2$ and X is **irreducible** otherwise.

A **hyperplane** $h \subseteq X$ is a non-empty connected subspace of X whose intersection with each n -cube σ of X is either empty or a midcube of σ (the space obtained from $[0, 1]^n$ by restricting one coordinate to $1/2$). It was proved in [Sag95] that each hyperplane of X separates X into two connected components, called **halfspaces**; we denote these h^+ and h^- . Given two subspaces $U, V \subseteq X$, we say that h **separates** U from V if U and V lie in distinct connected components of $X \setminus h$. Given two hyperplanes $h, k \subseteq X$, we say that h and k are **strongly separated** if no hyperplane in X intersects both h and k .

Let d denote the piecewise ℓ^1 -metric on X . The distance between any two vertices of X is precisely the number of hyperplanes separating them; this coincides with the restriction to the 0-skeleton of the path metric on the 1-skeleton of X . There is a well-defined **gate map** $\mathfrak{g}_h : X \rightarrow h$ which maps each vertex of X to its (unique) closest point projection in h (see [BHS17, Section 2.1]). This map has the property that *if x, y are vertices of X and k is a hyperplane then k separates $\mathfrak{g}_h(x)$ and $\mathfrak{g}_h(y)$ if and only if k intersects h and separates x from y* . In particular, if h and k are strongly separated hyperplanes then $\mathfrak{g}_h(k)$ is a single point.

We consider actions by cellular automorphisms, these are isometries with respect to both d and the CAT(0) metric (i.e. the piecewise ℓ^2 -metric). The group of all cellular automorphisms of X is denoted by $\text{Aut}(X)$. An action $G \curvearrowright X$ on a CAT(0) cube complex is said to be **proper** if the stabiliser of each n -cube is finite. It is **essential** if for any orbit $G \cdot v$ and for each halfspace h^+ bounded by a hyperplane h , there are points in $(G \cdot v) \cap h^+$ arbitrarily far from h .

We recall the Double Skewering Lemma of [CS11]: *Suppose that G is a group acting essentially on a finite dimensional CAT(0) cube complex without a fixed point at infinity. Then for any halfspaces $h^+ \subseteq k^+$, there is an element $\gamma \in G$ such that $\gamma k^+ \subsetneq h^+ \subseteq k^+$.*

Suppose that $h \subseteq X$ is a hyperplane such that the set of hyperplanes $G \cdot h$ is pairwise disjoint (i.e. for all $g \in G$ either $gh = h$ or $gh \cap h = \emptyset$) and such that G does not invert h (i.e. if $g \in \text{Stab}_G(h)$ then $gh^+ = h^+$). Let $N(h)$ be the open carrier of h ; that is, the union of all open cubes which intersect h . Let T_h be the graph whose vertex set is the set of connected components of $X \setminus (G \cdot N(h))$ and where $v, w \in V(T)$ are connected by an edge if they are both bounded by a common hyperplane carrier in $G \cdot N(h)$. This defines a continuous surjection $\pi_h : X \rightarrow T_h$ which maps each connected component of $X \setminus (G \cdot N(h))$ to a vertex and each open hyperplane carrier $N(k)$ in $G \cdot N(h)$ to an open edge whose midpoint is $\pi_h(k)$. Note that edges of T_h separate the space into two connected components so T_h is a tree. We will refer to T_h as the **dual tree** of h . The action $G \curvearrowright (G \cdot N(h))$ induces an action $G \curvearrowright T_h$ via π_h . Since G does not invert hyperplanes in the orbit of h , this action preserves the orientation of the edges, and is therefore an “action on a graph” in the sense of Section 5.

Recall that if $G \curvearrowright X$ is co-special then G acts with pairwise disjoint hyperplane orbits and no hyperplane inversions. By the above, this implies that G acts on the dual tree T_h for any hyperplane h of X .

Lemma 8.5. *Let G be a countable group acting on an irreducible CAT(0) cube complex X , where X is not quasi-isometric to \mathbb{R} and the automorphism group $\text{Aut}(X)$ acts cocompactly. Suppose that the action is proper, essential and does not fix a point in the visual boundary $\partial_\infty X$. Suppose moreover that there is a hyperplane $h \subseteq X$ such that the set of hyperplanes $G \cdot h$ is pairwise disjoint and such that G does not invert h . Let T_h be the corresponding dual tree. Under these conditions, the action $G \curvearrowright T_h$ is minimal and irreducible and there are edges $f_1, f_2 \in E(T)$ such that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite.*

Proof. Since the action of $\text{Aut}(X)$ is cocompact, X is finite dimensional.

We first show that there is a pair of strongly separated hyperplanes in the orbit $G \cdot h$. Since X is irreducible and finite dimensional and since G acts essentially without a fixed point at infinity, there is a pair of strongly separated hyperplanes $k_1, k_2 \subseteq X$ by [CS11, Theorem 5.1]. Since h does not intersect both k_1 and k_2 , then up to relabelling, we can assume that $h^- \subseteq k_2^-$ and that $k_2^+ \subseteq k_1^+$ where k_i^+, k_i^- are the halfspaces of k_i . By the Double Skewering Lemma, there exists $\gamma, \psi \in G$, such that $\gamma k_2^- \subsetneq k_1^-$ and $\psi h^+ \subsetneq k_2^+$. Thus $\gamma h \subseteq k_1^-$ and $\psi h \subseteq k_2^+$. If a hyperplane intersects both γh and ψh then it must intersect k_1 and k_2 so no such hyperplane exists. Therefore $h_1 := \gamma h$ and $h_2 := \psi h$ are strongly separated.

Recall that the image $\mathfrak{g}_{h_1}(h_2)$ is a single point, say x_1 . If $g \in \text{Stab}_G(h_1) \cap \text{Stab}_G(h_2)$ then $d(gx_1, h_2) = d(x_1, h_2)$ and, since $gx_1 \in h_1$ and x_1 is the unique point in $\mathfrak{g}_{h_1}(h_2)$, we have $gx_1 = x_1$. Since the action of G on X is proper, this implies that $\text{Stab}_G(h_1) \cap \text{Stab}_G(h_2)$ is finite. Thus if f_i is the edge of T_h with midpoint $\pi_h(h_i)$ for $i = 1, 2$ then $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite, as required.

Let us now show that the action $G \curvearrowright T_h$ is minimal and irreducible. Firstly, by the Double Skewering Lemma, T_h is unbounded and the G -action has no global fixed point. Since the action has a single orbit of edges, it is minimal.

Since X is irreducible and not a quasi-line and since the action of $\text{Aut}(X)$ is cocompact, [CS11, Lemma 7.1] implies that $\text{Aut}(X)$ does not stabilise any Euclidean flat \mathbb{R}^n in X . By [CS11, Theorem 7.2], this implies that there is a **facing triple** of hyperplanes in X : i.e. a

pairwise disjoint triple of hyperplanes k_0, k_1, k_2 such that no k_j separates the remaining two hyperplanes. Suppose that for each j , k_j^+ is the halfspace containing the remaining k_ℓ 's. Since h_1, h_2 are strongly separated, there is $i \in \{1, 2\}$ and $j \in \mathbb{Z}/3\mathbb{Z}$ such that $h_i \cap k_j = \emptyset$. By the Double Skewering Lemma, there exists $g \in G$ such that $gh_i \subseteq k_j^-$ (set $g = \text{id}$ if this was already the case). Then, using the Double Skewering Lemma again, there are $g_{j+1}, g_{j+2} \in G$ such that $g_{j+1}gh_i \subseteq k_{j+1}^-$ and $g_{j+2}gh_i \subseteq k_{j+2}^-$. The set $\{gh_i, g_{j+1}gh_i, g_{j+2}gh_i\}$ is a facing triple in the G -orbit of h . Thus there is a facing triple of hyperplanes (in this case mid-points of edges) in T_h , so T_h cannot be a line. By minimality of the action, G cannot fix a line in T_h .

It remains to show that G does not fix an end. Let $(e_n)_{n \in \mathbb{N}}$ be a geodesic ray such that e_1 is the edge whose midpoint is $\pi_h(h)$. There is a halfspace of h which contains $\pi_h^{-1}(e_n)$ for all $n \geq 2$, suppose without loss of generality that this halfspace is h^+ . By the Flipping Lemma of [CS11] there is an element $\gamma \in G$ such that $\gamma h^+ \subsetneq h^-$. Thus the paths $(e_n)_{n \geq 2}$ and $(\gamma e_n)_{n \geq 2}$ are contained in distinct components of $T_h \setminus \{e_1\}$ and therefore represent distinct ends of T_h . \square

Theorem 8.6. *If G is an infinite virtually compact special group then $G \in \mathcal{A}$.*

Examples of virtually compact special groups include right-angled Artin groups (essentially by definition); right-angled and hyperbolic Coxeter groups [NR03, HW10]; limit groups [Wis21, Theorem 18.6]; one-relator groups with torsion [Wis21, Corollary 19.2]. Many groups are known to act properly and cocompactly on CAT(0) cube complexes. By [Ago13], those that are moreover hyperbolic are then virtually compact special. This includes $C'(1/6)$ small cancellation groups [Wis04]; random groups at density $d < 1/6$ [OW11], hyperbolic-by-cyclic groups which are themselves hyperbolic [DMM23].

Proof. Recall that class \mathcal{A} contains every infinite amenable group and is closed under taking direct products. Moreover if G is virtually in \mathcal{A} then G itself is in \mathcal{A} . Thus we can assume that G is compact special. Let X be a CAT(0) cube complex on which G acts freely, cocompactly and co-specially. In particular, G is torsion-free. By [CS11, Proposition 3.5], there is a convex subcomplex of X , called the **essential core**, on which G acts essentially. The action of G on this subcomplex is free and cocompact and co-special. Thus by replacing X with its essential core and G with a finite index subgroup, we can assume that the action $G \curvearrowright X$ is free, cocompact, co-special and essential.

If X splits non-trivially as a product $X = X_1 \times X_2$ then by [Zal23, Lemma 3.8] there is a finite index subgroup $G' \leq G$ which splits as a direct product $G' = G_1 \times G_2$, where $G_1 = \text{Stab}_{G'}(X_1 \times \{x_2\})$ and $G_2 = \text{Stab}_{G'}(\{x_1\} \times X_2)$ for any $x_i \in X_i$. The actions $G_i \curvearrowright X_i$ are free, cocompact and co-special. Moreover, for each i , the dimension of X_i is strictly inferior to that of X .

By induction on the dimension, if we iterate the process of passing to the essential core, then applying [Zal23, Lemma 3.8] if the subspace is reducible, we arrive at a set of subgroups H_1, \dots, H_k of G such that:

- G can be constructed from H_1, \dots, H_k by taking direct products and finite index supergroups and
- each H_i acts on an irreducible CAT(0) cube complex X_i such that the action is free, cocompact, co-special and essential.

We can therefore assume that X is irreducible and $G \curvearrowright X$ has the above properties.

Let $\rho : X \rightarrow X \backslash G$ be the quotient map. Fix a hyperplane $h \subseteq X$ and let T_h be the corresponding dual tree. By [Wis21, Lemma 14.6] we can replace G with a finite index subgroup so that $\rho(h)$ does not separate $G \backslash X$. This implies that $G \backslash T_h$ is a loop.

By [CS11, Corollary 4.9], and since X is irreducible, either X is quasi-isometric to a line or there is no fixed point at infinity. In the former case, G is virtually cyclic (in fact, cyclic since it is torsion free) so $G \in \mathcal{A}$ since it is amenable. On the other hand, if G does not fix a point in $\partial_\infty X$ then it follows from Lemma 8.5 that the action of G on T_h is minimal and irreducible, and that there are edges $f_1, f_2 \in E(T_h)$ such that $\text{Stab}_G(f_1) \cap \text{Stab}_G(f_2)$ is finite. Since G is torsion-free this intersection is in fact trivial. Thus $G \in \mathcal{A}$ by Corollary 8.3. \square

A Appendix: A list of examples

A list of examples of finitely generated groups for which $\mathcal{K}(G) = \text{Sub}_{[\infty]}(G)$, thus $\text{rk}_{\text{CB}}(G) = 1$, and for which $G \curvearrowright \mathcal{K}(G)$ is highly topologically transitive.

1. Non-abelian limit groups (Corollary 5.23 and Example 5.22).
2. Non-elementary free products $A * B$ and groups with infinitely many ends and no non-trivial finite normal subgroup (Corollary 5.17). As for $\text{SL}(2, \mathbb{Z})$, see Example 5.18.
3. Graphs of free groups with cyclic edges such that at least one vertex group is non-cyclic (Corollary 5.23 and Example 5.22).
4. Groups in the class \mathcal{Q}' (Corollary 5.23).
5. Non-elementary and locally quasiconvex hyperbolic groups with trivial finite normal subgroup (Corollary 6.3). For instance, non-elementary hyperbolic groups with a small hierarchy and trivial finite normal subgroup (Example 6.5).
6. One-relator group $G = \langle S \mid w^m \rangle$ where w is a cyclically reduced word in $S \cup S^{-1}$ and $m \geq |w|_S$ (Example 6.6).
7. Coxeter groups $G = \langle s_1, \dots, s_n \mid s_i^2, (s_i s_j)^{m_{i,j}} \forall i < j \rangle$ where $\infty \geq m_{i,j} > n \geq 3$ (Example 6.7).

References

- [AFW15] Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton. *3-manifold groups*. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2015. 41
- [Ago04] Ian Agol. Tameness of hyperbolic 3-manifolds, 2004. arXiv:math/0405568. 4, 41
- [Ago08] Ian Agol. Criteria for virtual fibering. *Journal of Topology*, 1(2):269–284, 2008. 18
- [Ago13] Ian Agol. The virtual Haken conjecture. *Doc. Math.*, 18:1045–1087, 2013. With an appendix by Ian Agol, Daniel Groves, and Jason Manning. 4, 42, 48
- [Arz01] Goulnara N. Arzhantseva. On quasiconvex subgroups of word hyperbolic groups. *Geom. Dedicata*, 87(1-3):191–208, 2001. 37

- [BBS84] Andrew M. Brunner, Robert G. Burns, and Donald M. Solitar. The subgroup separability of free products of two free groups with cyclic amalgamation. In *Contributions to group theory*, volume 33 of *Contemp. Math.*, pages 90–115. Amer. Math. Soc., Providence, RI, 1984. [18](#)
- [Ben00] Ivar Bendixson. Quelques théorèmes de la théorie des ensembles de points, extrait d’une lettre adressée à M. Cantor à Halle. *Acta Mathematica*, 2:415 – 429, 1900. [8](#)
- [BF92] Mladen Bestvina and Mark Feighn. A combination theorem for negatively curved groups. *J. Differential Geom.*, 35(1):85–101, 1992. [36](#)
- [BFG23] Sasha Bontemps, Joshua Frisch, and Damien Gaboriau. The Cantor-Bendixson rank of the Baumslag group is infinite. Master’s thesis, 2023. [43](#)
- [BGK15] Lewis Bowen, Rostislav Grigorchuk, and Rostyslav Kravchenko. Invariant random subgroups of lamplighter groups. *Israel J. Math.*, 207(2):763–782, 2015. [3](#)
- [BHMS09] Martin R. Bridson, James Howie, Charles F. III Miller, and Hamish Short. Subgroups of direct products of limit groups. *Ann. Math. (2)*, 170(3):1447–1467, 2009. [32](#)
- [BHS17] Jason Behrstock, Mark F. Hagen, and Alessandro Sisto. Hierarchically hyperbolic spaces, I: Curve complexes for cubical groups. *Geom. Topol.*, 21(3):1731–1804, 2017. [46](#)
- [Bow12] Brian H. Bowditch. Relatively hyperbolic groups. *Internat. J. Algebra Comput.*, 22(3):1250016, 66, 2012. [37](#)
- [BW13] Hadi Bigdely and Daniel T. Wise. Quasiconvexity and relatively hyperbolic groups that split. *Michigan Math. J.*, 62(2):387–406, 2013. [36](#)
- [Can84] Georg Cantor. Über unendliche, lineare punktmannichfaltigkeiten, part 6. *Math. Ann.*, 23(4):453–488, 1884. [8](#)
- [Can96] Richard Canary. A covering theorem for hyperbolic 3-manifolds and its applications. *Topology*, 35(3):751–778, 1996. [4](#), [41](#)
- [CG86] Jeff Cheeger and Mikhael Gromov. L_2 -cohomology and group cohomology. *Topology*, 25(2):189–215, 1986. [17](#), [18](#), [31](#)
- [CG05] Christophe Champetier and Vincent Guirardel. Limit groups as limits of free groups. *Israel J. Math.*, 146:1–75, 2005. [5](#)
- [CG06] Danny Calegari and David Gabai. Shrinkwrapping and the taming of hyperbolic 3-manifolds. *Journal of the American Mathematical Society*, 19(2):385–446, 2006. [4](#), [41](#)
- [CGLM20] Alessandro Carderi, Damien Gaboriau, and François Le Maître. On dense totipotent free subgroups in full groups. *To appear in Geometry & Topology*, 2020. arXiv:2009.03080 [math.GR]. [3](#), [18](#)
- [CGLMS22] Alessandro Carderi, Damien Gaboriau, François Le Maître, and Yves Stalder. On the space of subgroups of Baumslag-Solitar groups I: perfect kernel and phenotype, 2022. arXiv:2210.14990 [math.GR]. [3](#), [36](#)

- [CGP10] Yves de Cornulier, Luc Guyot, and Wolfgang Pitsch. The space of subgroups of an abelian group. *Journal of the London Mathematical Society*, 81(3):727–746, 2010. [3](#)
- [CM87] Marc Culler and John W. Morgan. Group actions on \mathbb{R} -trees. *Proc. London Math. Soc. (3)*, 55(3):571–604, 1987. [21](#)
- [Coh89] Daniel E. Cohen. *Combinatorial group theory: a topological approach*, volume 14 of *Lond. Math. Soc. Stud. Texts*. Cambridge etc.: Cambridge University Press, 1989. [34](#)
- [Coo89] Michel Coornaert. Sur le domaine de discontinuité pour les groupes d’isométries d’un espace métrique hyperbolique. *Rend. Sem. Fac. Sci. Univ. Cagliari*, 59(2):185–195, 1989. [38](#)
- [Cor11] Yves Cornulier. On the Cantor-Bendixson rank of metabelian groups. *Annales de l’Institut Fourier*, 61(2):593–618, 2011. [8](#)
- [CS11] Pierre-Emmanuel Caprace and Michah Sageev. Rank rigidity for CAT(0) cube complexes. *Geom. Funct. Anal.*, 21(4):851–891, 2011. [47](#), [48](#), [49](#)
- [Dah03] François Dahmani. Combination of convergence groups. *Geom. Topol.*, 7:933–963, 2003. [36](#)
- [DK18] Cornelia Druţu and Michael Kapovich. *Geometric group theory*, volume 63 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2018. With an appendix by Bogdan Nica. [38](#)
- [DMM23] François Dahmani, Suraj Krishna MS and Jean Pierre Mutanguha. Hyperbolic hyperbolic-by-cyclic groups are cubulable, 2023. arXiv:2306.15054 [math.GR]. [48](#)
- [Fim14] Pierre Fima. Amenable, transitive and faithful actions of groups acting on trees. *Ann. Inst. Fourier (Grenoble)*, 64(1):1–17, 2014. [6](#)
- [GH90] Étienne Ghys and Pierre de la Harpe, editors. *Sur les groupes hyperboliques d’après Mikhael Gromov*, volume 83 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1990. Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988. [37](#), [38](#)
- [Git99] Rita Gitik. Ping-pong on negatively curved groups. *J. Algebra*, 217(1):65–72, 1999. [37](#)
- [GKM16] Yair Glasner, Daniel Kitroser, and Julien Melleray. From isolated subgroups to generic permutation representations. *J. London Math. Soc. (2)*, 94(3):688–708, 2016. [4](#), [17](#)
- [GM07] Y. Glasner and N. Monod. Amenable actions, free products and a fixed point property. *Bull. Lond. Math. Soc.*, 39(1):138–150, 2007. [6](#), [44](#)
- [Gro87] Mikhael Gromov. Hyperbolic groups. In S. M. Gersten, editor, *Essays in Group Theory*, pages 75–263. Springer New York, New York, NY, 1987. [3](#), [35](#), [37](#), [38](#)
- [Hal49] Marshall Hall, Jr. Subgroups of finite index in free groups. *Canad. J. Math.*, 1:187–190, 1949. [4](#), [17](#), [18](#)
- [HW01] G. Christopher Hruska and Daniel T. Wise. Towers, ladders and the B. B. Newman spelling theorem. *J. Aust. Math. Soc.*, 71(1):53–69, 2001. [36](#)

- [HW08] Frédéric Haglund and Daniel T. Wise. Special cube complexes. *Geom. Funct. Anal.*, 17(5):1551–1620, 2008. [4](#), [43](#), [46](#)
- [HW10] Frédéric Haglund and Daniel T. Wise. Coxeter groups are virtually special. *Adv. Math.*, 224(5):1890–1903, 2010. [48](#)
- [HW14] Pierre de la Harpe and Claude Weber. Malnormal subgroups and Frobenius groups: basics and examples. *Confluentes Math.*, 6(1):65–76, 2014. With an appendix by Denis Osin. [30](#)
- [Kec95] Alexander S. Kechris. *Classical Descriptive Set Theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag New York, NY, 1995. [8](#)
- [Kie20] Dawid Kielak. Residually finite rationally solvable groups and virtual fibring. *J. Amer. Math. Soc.*, 33:451–486, 2020. [18](#)
- [LS77] Roger C. Lyndon and Paul E. Schupp. *Combinatorial group theory*. Springer-Verlag, Berlin, 1977. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89*. [22](#)
- [Moo10] Soyoung Moon. Amenable actions of amalgamated free products. *Groups Geom. Dyn.*, 4(2):309–332, 2010. [6](#)
- [Moo11a] Soyoung Moon. Amenable actions of amalgamated free products of free groups over a cyclic subgroup and generic property. *Ann. Math. Blaise Pascal*, 18(2):211–229, 2011. [6](#), [45](#)
- [Moo11b] Soyoung Moon. Permanence properties of amenable, transitive and faithful actions. *Bull. Belg. Math. Soc. Simon Stevin*, 18(2):287–296, 2011. [6](#)
- [Mou88] Gabor Moussong. *Hyperbolic Coxeter groups*. ProQuest LLC, Ann Arbor, MI, 1988. Thesis (Ph.D.)—The Ohio State University. [36](#)
- [MP09] Eduardo Martínez-Pedroza. Combination of quasiconvex subgroups of relatively hyperbolic groups. *Groups Geom. Dyn.*, 3(2):317–342, 2009. [37](#)
- [MP12] Eduardo Martínez-Pedroza. On quasiconvexity and relatively hyperbolic structures on groups. *Geom. Dedicata*, 157:269–290, 2012. [37](#)
- [MT94] Michael L. Mihalik and Williams Towle. Quasiconvex subgroups of negatively curved groups. *J. Pure Appl. Algebra*, 95(3):297–301, 1994. [44](#)
- [MW08] Jonathan P. McCammond and Daniel T. Wise. Locally quasiconvex small-cancellation groups. *Trans. Amer. Math. Soc.*, 360(1):237–271, 2008. [36](#)
- [New68] Bill B. Newman. Some results on one-relator groups. *Bull. Amer. Math. Soc.*, 74:568–571, 1968. [36](#)
- [NR03] G.A. Niblo and L.D. Reeves. Coxeter groups act on CAT(0) cube complexes. *J. Group Theory*, 6(3):399–413, 2003. [48](#)
- [OW11] Yann Ollivier and Daniel T. Wise. Cubulating random groups at density less than $1/6$. *Trans. Amer. Math. Soc.*, 363(9):4701–4733, 2011. [48](#)
- [Pic06] Mikaël Pichot. Semi-continuity of the first ℓ^2 -Betti number on the space of finitely generated groups. *Comment. Math. Helv.*, 81(3):643–652, 2006. [18](#)
- [PT11] Jesse Peterson and Andreas Thom. Group cocycles and the ring of affiliated operators. *Invent. Math.*, 185(3):561–592, 2011. [31](#)

- [PV23] Luis Paris and Olga Varghese. Narrow normal subgroups of Coxeter groups and of automorphism groups of Coxeter groups, 2023. arXiv:2211.16906 [math.GR]. [36](#)
- [Rip82] Eliyahu Rips. Subgroups of small cancellation groups. *Bull. London Math. Soc.*, 14(1):45–47, 1982. [4](#), [43](#)
- [Sag95] Michah Sageev. Ends of group pairs and non-positively curved cube complexes. *Proc. London Math. Soc. (3)*, 71(3):585–617, 1995. [46](#)
- [Sco78] Peter Scott. Subgroups of surface groups are almost geometric. *J. London Math. Soc. (2)*, 17(3):555–565, 1978. [4](#), [17](#)
- [Ser80] Jean-Pierre Serre. *Trees*. Springer-Verlag, Berlin-New York, 1980. Translated from the French by John Stillwell. [19](#), [20](#), [33](#)
- [Sta68] John R. Stallings. On torsion-free groups with infinitely many ends. *Ann. of Math. (2)*, 88:312–334, 1968. [5](#), [31](#)
- [Sta71] John R. Stallings. *Group theory and three-dimensional manifolds*, volume 4 of *Yale Mathematical Monographs*. Yale University Press, New Haven, Conn.-London, 1971. A James K. Whittemore Lecture in Mathematics given at Yale University, 1969. [5](#), [31](#)
- [SW20] Rachel Skipper and Phillip Wesolek. On the Cantor-Bendixson rank of the Grigorchuk group and the Gupta-Sidki 3 group. *J. Algebra*, 555:386–405, 2020. [3](#)
- [Šir76] V. L. Širvanjan. Imbedding of the group $B(\infty, n)$ in the group $B(2, n)$. *Izv. Akad. Nauk SSSR Ser. Mat.*, 40(1):190–208, 223, 1976. [12](#)
- [Wil08] Henry Wilton. Hall’s theorem for limit groups. *Geom. Funct. Anal.*, 18(1):271–303, 2008. [18](#)
- [Wis04] D. T. Wise. Cubulating small cancellation groups. *Geom. Funct. Anal.*, 14(1):150–214, 2004. [48](#)
- [Wis21] Daniel T. Wise. *The structure of groups with a quasiconvex hierarchy*, volume 209 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2021. [46](#), [48](#), [49](#)
- [Zal23] Abdul Zalloum. Effective flipping, skewering and rank rigidity for cubulated groups with factor systems, 2023. arXiv:2305.16298 [math.GT]. [48](#)
- [Zel11] Yevhen G. Zelenyuk. *Ultrafilters and topologies on groups*, volume 50 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, 2011. [12](#)

P.A., UNIVERSITY OF BRISTOL, SCHOOL OF MATHEMATICS, BRISTOL, UK
 penelope.azuelos@bristol.ac.uk

D.G., CNRS, ENS-LYON, UNITÉ DE MATHÉMATIQUES PURES ET APPLIQUÉES, 69007 LYON, FRANCE
 damien.gaboriau@ens-lyon.fr